

Capture Of Toxic Pollutants By *Pistacia lentiscus* Leaves As A Low-Cost Biosorbent : Equilibrium, Kinetics And Thermodynamic Studies

Zerhouni Anissa, Bestani Benaouda*, Attouti Salima and Benderdouche Nouredine

Abdelhamid Ibn Badis University, Laboratory of Structure, Elaboration and Application of Molecular Materials (SEAMM), Faculty of Science and Technology, Mostaganem, Algeria

*Corresponding author, Email: bestanib@yahoo.fr; mel_anissa@yahoo.fr

Presenting expensive disposal problems during extraction of its essential oil, *Pistacia lentiscus* leaf was chosen in this investigation as a biosorbent for Rhodamine B and Pb^{2+} ions removal from simulated solution. Chemical and physico-chemical methods, such as FTIR analyses, minimum expenditure basket (MEB), mesoporous and microporous available areas and zero point charge (pH_{zpc}) were performed to characterize the biosorbent prior to its utilization. Effect of conventional parameters on biosorption of both pollutants, such as equilibrium time, adsorbent dose, pH and temperature were studied. Well known adsorption isotherms, namely Langmuir, Freundlich and Tempkin were used for adsorption equilibrium data analysis in their linear and non-linear forms. The rate of adsorption was encouraging for *P. lentiscus* waste used as such. Linearized and non-linearized Freundlich-biosorption models are more representative for the experimental data predicting heterogeneous surface coverage of the adsorbents. Magnitudes of R_L and n depict the favourability biosorption processes. Biosorption mechanism found to obey pseudo-second-order kinetic model and indicates that the sorption process is controlled by intra-particle diffusion. Thermodynamic analysis of the adsorption processes of both pollutants confirms their spontaneity and exothermicity. Compared to other biosorbents, *Pistacia lentiscus* leaves as a bio-renewable and affordable biomaterial can be efficiently used in removing organic and inorganic pollutants from industrial effluents.

KEYWORDS

Biosorption, *Pistacia lentiscus*, Isotherm models, Thermodynamics

REFERENCES

1. Krishna, I.V.M. and V. Manickam. 2017. Environmental management- Science and engineering for industry (1st edn). Butterworth-Heinemann.
2. Saifullahi, I. and F.B. Halimah. 2020. A short review on the removal of Rhodamine B dye using agricultural waste-based adsorbents. *Asian J. Chem. Sci.*, 7(1):25-37.
3. Rodriguez-Ramos, F.J. and E.R. Tonic. 2011. Use of Rhodamine B as a biomarker for oral plague vaccination of Prairie dogs. *J. Wildlife Diseases*. 47(3):765-768.
4. Bashir, W.B., et al. 2014. Application of solutions of Rhodamine B in dosimetry. *Appl. Radiation Isotopes*. 89:13-17.
5. Nestmann, E.R., et al. 1979. Mutagenic activity of Rhodamine dyes and their impurities as detected by mutation induction in Salmonella and DNA damage in Chinese hamster ovary cells. *Cancer Res.*, 39(11):4412-4417.
6. Sigma-Aldrich. 2015. Rhodamine B (Material Safety Data Sheet) Version 5. 4. Available at: <http://www.sigmaaldrich.com/MSDS/DisplayMSDSPage.do?country=BN&language=en&productNumber=R4127&brand=SIGMA> & Page. To Go To URL=[http%3A%2F%2Fwww.sigmaaldrich.com%2Fcatalog%2Fproduct%2Fsigma%2Fr4127%3Flang%30en1](http://www.sigmaaldrich.com/catalog/product/sigma/r4127).
7. Wani, A., A.R.A. Anjum and A.J. Usmani. 2015. Lead toxicity : A review. *Interdiscip. Toxicol.*, 8(2): 55-64.
8. ATSDR. 2012. Case studies in environmental medicine-Lead (Pb) toxicity. Agency for Toxic Substances and Disease Registry.
9. Gagan, F., D. Gupta and A. Tiwari. 2012. Toxicity of lead : A review with recent updates. *Interdiscip. Toxicol.*, 5(2): 47-58.

10. Grant, L.D. 2020. Lead and compounds. In Environmental toxicants : Human exposures and their health effects (4th edn). Ed M. Lippmann and G.D. Leikauf. John Wiley & Sons Inc.
11. Street, A. and W.O. Alexander. 1998. Metals in the service of man (11th edn). Penguin, UK.
12. Rui, Z., et al. 2015. Source of lead pollution, its influence on public health and the counter measures. *Int. J. Health. Animal Sci. Food Safety.* 2: 18-31.
13. Bakala'r, T., M. Bugel and L. Gajosova. 2009. Heavy metal removal using reverse osmosis. *Acta Montanistica Slovaca.* 14(3):250-253.
14. Yossor, R.A. and H.A. Ahmed. 2016. Removal of heavy metals from industrial wastewater by using RO membrane. *Iraqi J. Chem. Petroleum Eng.,* 17(4): 125-136.
15. Gaikwad, R.W., V.S. Sapkal and R.S. Sapkal. 2010. Ion exchange system design for removal of heavy metals from acid mine drainage wastewater. *Acta Montanistica Slovaca Rocknik.* 4: 298-304.
16. Benderdouche, N., et al. 2003. Enhancement of the adsorptive properties of a desert *Salsola vermiculata* species. *Adsorption Sci. Tech.,* 21(8).
17. Azimi, A., et al. 2017. Removal of heavy metals from industrial wastewaters : A review. *Chem. Bio. Eng. Rev.,* 4(1): 1-24.
18. Fomina, M. and G.M. Gadd. 2014. Biosorption : Current perspectives on concepts, definition and application. *Bioresour. Tech.,* 160:3-14.
19. Arshid, B., et al. 2019. Removal of heavy metal ions from aqueous system by ion-exchange and biosorption methods. *Env. Chem. Letters.* 17: 729-754.
20. Michalak, J., K. Chojnacka and W. Krowiak. 2013. A state of the art for the biosorption process - A review. *Appl. Biochem. Biotech.,* 170 (6): 1389-1416.
21. Agarwal, R. and M.K. Singh. 2017. Heavy metal removal from wastewater using various adsorbents : A review. *J. Water Reuse Desal.,* 7(4): 387-419.
22. Ouldoumna, A., et al. 2013. Characterization and application of three novel biosorbents '*Eucalyptus globulus*, *Cynara cardunculus* and *Prunus cerasifera*' to dye removal. *Desal. Water Treatment.* 51: 3527-3538.
23. De Frietas, G.R., M.G.C. de Silva and M.G.A. Vieira. 2019. Biosorption technology for removal of toxic metals : A review of commercial biosorbents and patents. *Env. Sci. Poll. Res. Int.,* 26(19): 19097-19118.
24. Guiyin, W., et al. 2018. Removal of Pb (II) from aqueous solutions by *Phytolacca americana* L. biomass as a low cost biosorbent. *Arabian J. Chem.,* 11: 99-110.
25. Farnane, M., et al. 2018. New sustainable biosorbent based on recycled deoiled carob seeds : Optimization of heavy metals remediation. *J. Chem.* DOI:10.1155/2018/574 8493.
26. Haouli, A., et al. 2015. Contribution to the analysis of *Pistacia lentiscus* extracted oil. *American-Eurasian J. Agric. Env. Sci.,* 15(6): 1075-1081.
27. Tingshuang, Y., et al. 2008. Phylogenetics and reticulate evolution of *Pistacia* (Anacardiaceae). *American J. Botany.* 95(2):241-251.
28. Benyoussef, E.H., et al. 2005. Essential oil of *Pistacia lentiscus* L. from Algeria. *J. Essential Oil Res.,* 17: 642-644.
29. Bampouli, A., et al. 2014. Comparison of different extraction methods of *Pistacia lentiscus* var. chia leaves : Yield, antioxidant activity and essential oil chemical composition. *J. Appl. Res. Medicinal Aromatic Plants.* 1(3): 81-91.
30. Attoub, S., et al. 2014. Short-term effects of oral administration of *Pistacia lentiscus* oil on tissue-specific toxicity and drug metabolizing enzymes on mice. *Cell Physical. Biochem.,* 33(5): 1400-1410.
31. Haloui, T., et al. 2015. The use of *Pistacia lentiscus* L. oil as green inhibitor for corrosion of mild steel in 1 M hydrochloric acid solution : Thermodynamic and adsorption. *Der Pharma Chemica.* 7(9): 225-238.
32. Dahmoune, F., et al. 2014. *Pistacia lentiscus* leaves as a source of phenolic compounds : Microwave-assisted extraction optimized and compared with ultrasound assisted and conventional solvent extraction. *Ind. Crops Products.* 61: 31-40.
33. Cherbal, A., et al. 2012. Extraction and valorization of phenolic compounds of leaves of Algerian *Pistacia lentiscus*. *Asian J. Plant Sci.,* 11:131-136.
34. Cheurfa, M. and R. Allem. 2015. Study of hypo-cholesterolemic activity of Algerian *Pistacia lentiscus* leaves extracts in vivo. *Rev. Bras. Farmacogn.,* 25(2).
35. Haloui, T., et al. 2015. Effect of harvesting period and drying time on the essential oil yield of *Pistacia lentiscus* L. leaves. *Der Pharma Chemica.* 7(10): 320-324.

36. Bestani, B., *et al.* 2008. Adsorption of methylene blue and iodine from aqueous solutions by a desert *Salsola vermiculata* species. *Bioresour.*, 99(17): 8441-8444.
37. D 4607-94. 2006. Standard test method for determination of iodine number of activated carbon. ASTM International, U.S.
38. Kaewpravit, C., *et al.* 1998. Application of methylene blue adsorption to cotton fiber specific surface area measurement: Part I. Methodology. *J. Cotton Sci.*, 173: 2164.
39. Benzekri, M.B., *et al.* 2018. Valorization of olive stones into a granular activated carbon for the removal of methylene blue in batch and fixed bed modes. *J. Mater. Env. Sci.*, 9(1): 272-284.
40. Noszko, L.H., *et al.* 1984. Preparation of activated carbon from the byproducts of agricultural industry. *Per. Polytech.*, 28: 293-297.
41. Cleiton, A., C. Nunes and M. Guerreiro. 2011. Estimation of surface area and pore volume of activated carbons by methylene blue and iodine numbers. *Quim. Nova.*, 34: 472-476.
42. Ghasemi, M., *et al.* 2014. Adsorption of Pb (II) from kinetic studies. *J. Ind. Eng. Chem.*, 20: 2193-2199.
43. Yuvaraja, G., *et al.* 2014. Biosorption of Pb (II) from agricultural waste. *Colloids Surf.*, B114: 75-81.
44. Melichova, Z. and L. Hromada. 2013. Adsorption of Pb²⁺ and Cu²⁺ ions from aqueous solutions on natural bentonite. *Polish J. Env. Stud.*, 22(2): 457-464.
45. Kooh, M.R., *et al.* 2016. Separation of toxic Rhodamine B from aqueous solution using an efficient low-cost material, *Azolla pinnata* by adsorption method. *Env. Monit. Assess.*, 188(2): 108.
46. Freundlich, H.M.F. 1906. Adsorption in solutions. *J. Physical Chem.*, 57: 385-470.
47. Langmuir, I. 1916. The constitution and fundamental properties of solids and liquids. Part I. Solids. *J. American Chem. Soc.*, 38:2221-2295.
48. Tempkin, M.J. 1941. Adsorption equilibrium and the kinetics of process on non-homogenous surfaces and in the interaction between adsorbed moisture. *J. Phys. Chem. (USSR)*. 15: 296-332.
49. Svecova, L., *et al.* 2006. Cadmium, lead and mercury biosorption on waste fungal biomass issued from fermentation industry. I. Equilibrium studies. *Sep. Purif. Tech.*, 52: 142-153.
50. Xin, H., *et al.* Pb²⁺ biosorption from aqueous solutions by live and dead biosorbents of the hydrocarbon-degrading strain *Rhodococcus* sp. Hx-2. *PLoS One*. 15(1). DOI: 10.1371/journal.pone.0226557.
51. Santhi, T., A.L. Prasad and A. Manonmani. 2014. A comparative study of microwave and chemically treated *Acacia nilotica* leaf as an eco-friendly adsorbent for the removal of Rhodamine B dye from aqueous solution. *Arabian J. Chem.*, 7: 494-503.
52. Kooh, M.R., M.K. Dahri and L.B. Lim. 2016. Remediation of Rhodamine B dye from aqueous solution using *Casuarina equisetifolia* cone powder as a low-cost adsorbent. *Adv. Physical Chem.* DOI:10.1155/2016/9497378.
53. Lagergren, S. 1898. About the theory of so-called adsorption of soluble substances. *Royal Swedish Academy Sci.*, 24:1-39.
54. Ho, Y.S. and G. McKay. 1999. Pseudo second order model for sorption processes. *Process Biochem.*, 34(5): 451-465.
55. Bullen, J., S. Saleesongsom and D. Weiss. 2000. A revised pseudo second order kinetic model for adsorption, sensitivity to changes in sorbate and sorbent concentrations. *Physical Chem.* DOI: 10.26434/chemrxiv.12008799.v1.
56. Weber Jr., W.J. and J.C. Morris. 1963. Kinetics of adsorption on carbon from solution. *J. Sanitary Eng. Div. Proceed. American Soc. Civil Eng.*, 89: 31-59.
57. Vasanth, K.V., V. Ramamurthi and S. Sivannesan. 2005. Modeling the mechanism involved during the sorption of methylene blue onto flyash. *J. Colloid Interface Sci.*, 284: 14-21.

Determination Of Rare Earth Elements And Its Distribution Pattern From The Core Sediments By K_0 -Instrumental Neutron Activation Analysis

Ebenezer Aquisman Asare^{1,2*}, Rafeah Wahi¹, Alexander Obiri Gyampoh³, and Omolayo Ajoke Omorinoye⁴

1. Universiti Malaysia Sarawak, Resource Chemistry Programme, Faculty of Resource Science and Technology, Kota Samarahan, Sarawak, Malaysia

2. University of Ghana, Graduate School of Nuclear and Allied Sciences, Kwabenya-Accra, Ghana

3. Kibi Presbyterian College of Education, Departments of Science, Kibi, Eastern Region, Ghana

4. University of Ilorin, Faculty of Physical Sciences, Department of Geology and Mineral Sciences, Ilorin, Nigeria

*Corresponding author, Email : aquisman1989@gmail.com

This work aimed to assess rare earth elements and their distribution pattern from the core sediments from the central coast of Ghana by K_0 instrumental neutron activation analysis. The rare earth element content was evaluated with uncertainty less than 8% (at 95% confidence level) and demonstrated to be accordant with the IAEA-soil 7 certified concentrations. The calculated concentration of light rare earth elements and Fe normalized enrichment factors suggested that sediment samples were not enriched with light rare earth elements (LREEs) obtained from discharges of anthropogenic activities. The chondrite-normalized pattern of rare earth elements exhibited LREEs, Tm, Tb, Eu and Ho enrichment. The total contents of rare earth elements calculated can be used to establish baseline information about environmental contamination determination and to develop the relationships between the Ce/Ce* and Eu/Eu* anomalies and the source appointment of both LREEs elements and heavy rare earth elements (HREEs).

KEYWORDS

Rare earth element, Marine core sediment, Neutron activation analysis, Shale average, Enrichment factor

REFERENCES

- Adam, J.R., *et al.* 2003. Isotope dilution MC-ICP-MS isotope dilution MC-ICP-MS rare earth element analysis of geo-chemical reference materials NIST SRM 610, SRM 612, SRM 614, BHVO-2G, BHVO-2, BCR-2G, JB-2, WS-E, W-2, AGV-1 and AGV-2. *Geo Standard Geo Anal. Res.*, 28:417–429.
- Ozdes, D., C. Diran and B.H. Senturk. 2011. Adsorptive removal of Cd (II) and Pb (II) ions from aqueous solutions by using Turkish illitic clay. *J. Env. Manage.*, 92(12):3082–3090.
- Henderson, P. 1984. General geochemical properties and abundances of the rare earth elements. In *Rare earth element geochemistry* (vol. 2). pp 1-32.
- Khadijeh, R., *et al.* 2009. Rare earth elements determination and distribution patterns in surface marine sediments of the South China sea by INAA, Malaysia. *J. Rare Earths*. 27 (6):1066-1071.
- McLennan, S.M. 1989. Rare earth elements in sedimentary rocks. Influence of province and sedimentary processes In *Geochemistry and mineralogy of rare earth elements*.
- Shatrov, V. A., *et al.* 2008. Rare earth elements as indicators of the tectonic activity of the basement (with reference to the Voronezh antecline). *Doklady Earth Sci.*, 423(2):1467.
- Al-Zahrany, A.A. 2007. Elemental distribution of marine sediments in the Sbaitis of Malacca by utilizing neutron activation and mass spectro-scopy analysis, Ph.D Thesis. University Putra, Malaysia.
- Alnour, I. A., *et al.* 2014. Investigate the capability of INAA absolute method to determine the concentrations of ²³⁸U and ²³²Th in rock samples. *J. Radioanal. Nucl. Chem.*, 299:177-186.
- Kafala, S. and T. MacMahon. 2007. Comparison of neutron activation analysis methods. *J. Radioanal. Nucl. Chem.*, 271(2): 507-516.
- Duffield, J. and G. Gilmore. 1979. An optimum method for the determination of rare earth elements by neutron activation analysis. *J. Radioanal. Nucl. Chem.*, 48(1-2):35-145.

11. Danko, B., R. Dybczynski and Z. Samczynski. 2008. Accurate determination of individual lanthanides in biological materials by NAA with pre- and post-irradiation separation. *J. Radioanal. Nucl. Chem.*, 278(1):81-88.
12. Ravisankar, R., *et al.* 2006. Determination and distribution of rare earth elements in beach rock samples using instrumental neutron activation analysis (INAA). *Nuclear Instruments Methods Physics Res. Section B: Beam Interactions Mater. Atoms.*, 251(2):496-500.
13. Silachyov, I. 2016. Rare earths analysis of rock samples by instrumental neutron activation analysis, internal standard method. *J. Radioanal. Nucl. Chem.*, 310(2):573-582.
14. Abdollahi, N., *et al.* 2018. Novel approach in k_0 -NAA for highly concentrated REE samples. *Talanta*. 180:403-409.
15. Dampare, S. B., *et al.* 2005. Determination of rare earth elements by neutron activation analysis in altered ultramafic rocks from the Akwatia district of the Birim diamondiferous field, Ghana. *J. Radioanal. Nucl. Chem.*, 265(1):101-106.
16. Whitty-Leveille, L., *et al.* 2017. A comparative study of sample dissolution techniques and plasma-based instruments for the precise and accurate quantification of REEs in mineral matrices. *Anal. Chimica Acta*. 961:33-41.
17. Santoro, A., *et al.* 2016. Assessing rare earth elements in quartz rich geological samples. *Appl. Radiation Isotopes*. 107:323-329.
18. Zawisza, B., *et al.* 2011. Determination of rare earth elements by spectroscopic techniques: A review. *J. Anal. Atomic Spectrometry*. 26(12): 2373-2390.
19. Ravichandran, M., M. Baskaran and P.H. Santschi. 1995. History of trace metal pollution in Sabine-Neches Eshury, Beaumon, Texas. *Env. Sci. Tech.*, 29:1495.
20. Lide, D. R. 2005. Geophysics, astronomy and acoustics abundance of elements in the earth's crust and in the sea. Handbook of chemistry and physics (85th edn). CRC Press, Boca Raton, Florida.
21. Hannigan, R., E. Dorval and J. Cynthia. 2010. The rare earth element chemistry of estuarine surface sediments in the Chesapeake bay. *Chem. Geol.*, 272:20-30.

Distribution Of Indoor Air Pollutants Relative To Meteorological Parameters In Selected Schools In The Eastern Cape Province Of South Africa: A Preliminary Study

Anye Chungag^{1*}, Godwill A. Engwa² and Benedicta N. Nkeh-Chungag²

1. Dayenuel Consulting, Postnet Suites 092, Mthatha 5099. South Africa

2. Walter Sisulu University, Department of Biological and Environmental Sciences, Faculty of Natural Sciences, Mthatha, South Africa

*Corresponding author, Email : foyarrh52@gmail.com

Particulate matter (PM), gaseous pollutants and meteorological parameters have gained concern recently in indoor air quality (IAQ) assessment. However, information about the nature and concentration of indoor air pollutants and meteorological parameters in the presence and absence of learners is scarce, hence the aim of this study. This study sampled indoor air from 23 classrooms in three urban and four rural schools in the Eastern Cape Province of South Africa. Components of indoor air (O_2 , CO_2 , NO_2 , $PM_{2.5}$, PM_5 and PM_{10}) and meteorology (relative humidity, wet bulb temperature, dew point temperatures, barometric pressure and airflow) were measured. The findings showed that sampled pollutants were present in all the classrooms in different proportions and varied with the presence and/or absence of learners. While CO_2 and PM concentrations were higher than permissible amounts, NO_2 volumes were barely detectable. The meteorological parameters were comparatively much higher in urban than rural school classrooms in the presence and absence of learners. In conclusion, classrooms present different IAQ when learners are present compared to when they are absent considering the concentration of their ambient compromisers. The levels of air pollutants in school classrooms do not only increase with the presence of learners but tend to also increase with the rise in indoor meteorological factors.

KEYWORDS

Indoor air quality, Air pollution, Particulate matter, Gas pollutants, Meteorological parameters

REFERENCES

1. WHO. 2012. Guidelines for indoor air quality: Selected pollutants. World Health Organisation Regional Office for Europe, Copenhagen, Denmark.
2. Tang, T., *et al.* 2012. Fine and ultrafine particles emitted from laser printers as indoor air contaminants in German offices. *Env. Sci. Poll. Res.*, 19(9): 3840-3849.
3. Khaefi, M., *et al.* 2017. Association of particulate matter impact on prevalence of chronic obstructive pulmonary disease in Ahvaz, southwest Iran during 2009-2013. *Aerosol Air Quality Res.*, 17(1): 230-237.
4. Choi, S. H., *et al.* 2014. Behaviour of particulate matter during high concentration episodes in Seoul. *Env. Sci. Poll. Res.*, 21(9):5972-5982.
5. Maggos, T., *et al.* 2017. Development of standardized method on automated measuring systems for particulate matter PM_{10} , $PM_{2.5}$: Field validation tests. *Fresen. Env. Bull.*, 26(1): 273-282.
6. Goyal, S.K. and C.C. Rao. 2007. Assessment of atmospheric assimilation potential for industrial development in an urban environment: Kochi (India). *Sci. Total Env.*, 376 (1-3): 27-39.
7. Zhang, H., *et al.* 2015. Relationships between meteorological parameters and criteria air pollutants in three megacities in China. *Env. Res.*, 140: 242-254.
8. Griffiths, M. and M. Eftekhari. 2008. Control of CO_2 in a naturally ventilated classroom. *Energy Build.*, 40: 556-560.
9. Kankaria, A., B. Nongkynrih and K.S. Gupta. 2014. Indoor air pollution in India : Implications on health and its control. *J. Comm. Med.*, 39(4): 203-207.
10. Chatzidiakou, L., D. Mumovic and A.J. Summerfield. 2012. What do we know about indoor air quality in school classrooms? A critical review of the literature. *Intelligent Build. Int.*, 4(4): 228-259.

11. Ranjbar, A. and Y. Afacan. 2018. Analysing the effects of thermal comfort and indoor air quality in design studios and classrooms on student performance. Department of Interior Architecture and Environmental design, Bilkent University. Easy Chair Preprint. No. 435.
12. Chatzidiakou, L., D. Mumovic and J. Summerfield. 2015. Is CO₂ a good proxy for indoor air quality in classrooms? Part 1: The interrelationships between thermal conditions, CO₂ levels, ventilation rates and selected indoor pollutants. *Build. Services Eng. Resour. Tech.*, 36: 129–161.
13. Johna, J., *et al.* 2004. Our children in day care: Reducing exposure to environmental lead at day care centres. *South Africa J. Sci.*, 100: 135-138.
14. Versteeg, H. 2007. Research into the quality of the indoor environment in primary schools. The Hague : Ministry of Housing, Spatial Planning and the Environment.
15. Meijer, G. and F. Duijm. 2009. Indoor environment of the public schools in Groningen. GGD Groningen.
16. Habets, T., *et al.* 2008. Assessment of ventilation in schools. Utrecht : GGD Netherland.
17. WHO. 2011. Guidelines for indoor air quality. Selected pollutants. World Health Organisation, Geneva, Switzerland. pp 55-89. Available at : <http://www.ncbi.nlm.nih.gov/books/NBK138710/>.
18. Shendell, D.G., *et al.* 2004a. Associations between classroom CO₂ concentrations and student attendance in Washington and Idaho. *Indoor Air*. 14 (5): 333–341.
19. Bako-Biro, Z., *et al.* 2012. Ventilation rates in schools and pupils' performance. *Build. Env.*, 48: 215-223.
20. Blondeau, P., *et al.* 2005. Relationship between outdoor and indoor air quality in eight French schools. *Indoor Air*. 15 (1): 2–12.
21. Chaloulakou, A., I. Mavroidis and A. Duci. 2003. Indoor and outdoor carbon monoxide concentration relationships at different microenvironments in the Athens area. *Chemosphere*. 52 (6): 1007–1019.
22. Van Roosbroeck, S., *et al.* 2007. Long-term personal exposure to PM_{2.5}, soot and NO_x in children attending schools located near busy roads, a validation study. *Atmos. Env.*, 41(16): 3381–3394.
23. Nkonkobe Municipality. 2012. Alice regeneration programme: High level feasibility assessment report. Nkonkobe Municipality, Integrated Development Plan. pp 1-59.
24. SAAQIS. 2017. South African air quality information system. National priority area monthly ambient air quality monitoring station reports, Department of Environmental Affairs, South Africa.
25. Salthammer, T. 2011. Critical evaluation of approaches in setting indoor air quality guidelines and reference values. *Chemosphere*. 82: 1507–1517.
26. Schneider, M. 2002. Do school facilities affect academic outcomes? National clearing house for educational facilities, Washington D.C.
27. Petzer, G. 2010. Indoor air quality issues in South Africa. Airshed Planning Professionals, Halfway House, South Africa.
28. Kunkel, S. and E. Kontonasiou. 2016. Indoor air quality, thermal comfort and daylight policies on the way to nZEB – Status of selected MS and future policy recommendations. ECEEE Proceedings on Policies and programmes towards a zero-energy building stock.
29. Mendell, M. J., *et al.* 2013. Association of classroom ventilation with reduced illness absence: A prospective study in California elementary schools. *Indoor Air*. 23: 515–528.
30. Yang, W., *et al.* 2009. Indoor air quality investigation according to age of the school buildings in Korea. *J. Env. Manage.*, 90: 348–354.
31. Rovelli, S., *et al.* 2014. Airborne particulate matter in school classrooms in Northern Italy. *Int. J. Env. Res. Public Health*. 11(2):1398-1421.
32. Miller, G.T. 1998. Living in the environment: An introduction to environmental science. Wadsworth Publishers.
33. Morawska, L., *et al.* 2003. Characteristics of particle number and mass concentrations in residual houses in Brisbane. *Australia Atmos. Env.*, 37: 4195– 4203.
34. WHO. 2013. Health effects of particulate matter. Policy implications for countries in Eastern Europe, Caucasus and Central Asia. World Health Organization, Geneva.
35. Heger, M. and M. Sarraf. 2018. Air pollution in Tehran: Health costs, sources and policies. Environment and Natural Resources Global Practice Discussion Paper No. 6. World Bank, Washington DC.
36. Nunes, R. A. O., *et al.* 2015. Particulate matter in rural and urban nursery schools in Portugal. *Env. Poll.* DOI : 10.1016/j.envpol.2015.03.009.
37. Goyal, R. and M. Khare. 2009. Indoor – outdoor concentrations of RSPM in classroom of a naturally ventilated school building near an urban traffic roadway. *Atmos. Env.*, 43(38): 6026–6038.

38. Heudorf, U., V. Neitzert and J. Spark. 2009. Particulate matter and carbon dioxide in classrooms- The impact of cleaning and ventilation. *Int. J. Hyg. Env. Health.* 212(1):45-55.
39. Fromme, H., *et al.* 2007. Particulate matter in the indoor air of classrooms- Exploratory results from Munich and surrounding area. *Atmos. Env.*, 41: 854–866.
40. Heudorf, U., V. Neitzert and J. Spark. 2009. Particulate matter and carbon dioxide in classrooms- The impact of cleaning and ventilation. *Int. J. Hyg. Env. Health.* 212: 45-55.
41. Gold, D. R., *et al.* 1999. Particulate and ozone pollutant effects on the respiratory function of children in southwest Mexico City. *Epidemiol.*, 10(1): 8–16.
42. Peterson, J. 2017. The relationship between mois ture and temperature. *Sciencing.*
43. Fang, Z., *et al.* 2019. Investigation into the thermal comfort of university students conducting outdoor training. *Build. Env.*, 149:26-38.
44. Tucker, W.G. 1989. ASHRAE standard 62: ventilation for acceptable indoor air quality. U.S. Environmental Protection Agency.
45. ASHRAE. 2017. ASHRAE standard 55 - Thermal environmental conditions for human occupancy. ANSI/ABHRAE.
46. Wolkoff, P. and S. K. Kjaergaard. 2007. The dichotomy of relative humidity on indoor air quality. *Env. Int.*, 33(6): 850–857.

Mapping Of Aquifer Vulnerability Zones Using Drastic-Lu Model For The Part Of Palwal – Faridabad District, Yamuna Alluvial Aquifer, Haryana, India

Nepal Singh and Sarfaraz Ahmad*

Aligarh Muslim University, Department of Geology, Aligarh - 202 002, India

*Corresponding author, Email : sarf71@gmail.com; nepalsingh.amu@gmail.com

Vulnerability assessment to delineate areas that are more susceptible to contamination from anthropogenic source has become an important element for sensible resource management and landuse planning. The study utilized the DRASTIC-Lu model comprising seven environmental parameters which include depth to the water table, net recharge, aquifer media, soil media, topography, the impact of the vadose zone, hydraulic conductivity. These thematic maps under the GIS environment were used to assess the aquifer vulnerability in Yamuna alluvial aquifer in parts of Palwal - Faridabad district, Haryana, India. The result of the groundwater vulnerability assessment suggests four zones of relative vulnerability as very low, low, medium and highest vulnerable zones. The elevated north-western parts of the study area displayed high aquifer vulnerability due to shallow groundwater depth, open water bodies, urban settlement, soil type and high hydraulic conductivity. While eastern parts show less vulnerability due to low hydraulic conductivity and deeper groundwater table.

KEYWORDS

DRASTIC-Lu, Aquifer vulnerability mapping, Yamuna alluvial, Palwal - Faridabad, GIS

REFERENCES

1. Tesoriero, A.J., E.L. Inkpen and F.D. Voss. 1998. Assessing groundwater vulnerability using logistic regression. Source water assessment and protection 98, Technical Conference. Proceedings, pp 157-165.
2. Zhang, R., *et al.* 1996. Determination of non-point source pollution using GIS and numerical models. *J. Env. Quality*. 25: 411-418
3. Aller, L., *et al.* 1985. DRASTIC : A standardized system for evaluating groundwater pollution potential using hydrogeologic settings. USEPA Report 600/2-87/035. U.S. Environmental Protection Agency. Available at : <http://www.epa.gov/nscep>.
4. Dixon, B. 2005. Applicability of neuro-fuzzy techniques in predicting groundwater vulnerability : A GIS-based sensitivity analysis. *J. Hydrol.*, 309:17-38.
5. Foster, S.S.D. 1987. Fundamental concept in aquifer vulnerability pollution risk and protection strategy. In The Haque vulnerability of soil and ground- water to pollutants, proceeding and information. Ed Duijvenbooden W. Van and Waegeninng H.G. Van. TNO Committee on Hydrological Research. 38:69-86.
6. Worrall, F. and D.W. Kopin. 2004. Aquifer vulnerability to pesticide pollution-combining soil, Land-use and aquifer properties with molecular descriptors. *J. Hydrol.* 29: 191-204.
7. Kalinski, R.J., *et al.* 1994. Correlation between DRASTIC vulnerability and incidents of VOC contamination of municipal wells in Nebraska. *Ground Water*. 32(1):31-34.
8. Hussain, Y., *et al.* 2017. Groundwater quality evaluation by electrical resistivity method for optimized tubewell site selection in an arid stressed Thal Doab aquifer in Pakistan. *Modeling Earth Syst. Env.* DOI:10.1007/s40808-017-0282-3.
9. Al-Abadi, A.M., A.M. Al-Shamm'a and M.H. Aljabbari. 2014. A GIS-based DRASTIC model for assessing intrinsic groundwater vulnerability in northeastern Missan Governorate, Southern Iraq. *Appl. Water Sci.*, 7:89-10.
10. Shirazi, S.M., H.M. Imran and S. Akib. 2012. GIS-based DRASTIC method for groundwater vulnerability assessment : A review. *J. Risk Res.*, 15 (8):991-1011.

11. Thapinta, A. and P.F. Hudak. 2003. Use of geographic information systems for assessing ground-water pollution potential by pesticides in Central Thailand. *Env. Int.*, 29:87-93.
12. Umar, R., I. Ahmad and F. Alam. 2009. Mapping groundwater vulnerability zones using modified DRASTIC approach of an alluvial aquifer in parts of Central Ganga plain, Western Uttar Pradesh. *J. Geol. Soc. India*. 73:193-201.
13. Rehman, A. 2008. A GIS based DRASTIC model for assessing groundwater vulnerability in shallow aquifer in Aligarh, India. *Appl. Geo.*, 28(1): 32-53.
14. Khan, M., R. Umar and H. Lateh. 2010. Assessment of aquifer vulnerability in parts of Indo Gangatic plain. *Int. J. Phy. Sci.*, 5 (III):1711-1720.
15. Ghosh, A., A.K. Tiwari and S. Das. 2015. A GIS based DRASTIC model for assessing groundwater vulnerability of Katri watershed, Dhanbad, India. *Modeling Earth Sys. Env.* DOI:10.1007/S40808-015-0009-2.
16. Ghosh, T. and R. Kanchan. 2016. Aquifer vulnerability assessment in the Bengal alluvial tract, India, using GIS based DRASTIC model. *Model. Earth Sys. Env.* DOI:10.1007/S40808-016-0208-5.
17. Kumar, P., *et al.* 2016. Assessment of the effectiveness of DRASTIC in predicting the vulnerability of groundwater to contamination : A case study from Fatehgarh Sahib district in Punjab. *Env. Earth Sci.*, 75(10):1-13.
18. Prasad, K. and J.P. Shukla. 2014. Assessment of groundwater vulnerability using GIS based DRASTIC technology for the basaltic aquifer of Burhner watershed, Mohgaon block, Mandla (India). *Curr. Sci.*, 107(10):1649-1656.
19. Ahada, C.P.S. and S. Suthar. 2018. A GIS based DRASTIC model for assessing aquifer vulnerability in southern Punjab, India. *Modeling Earth Sys. Env.* DOI:10.1007/s40808-018-0449-6.
20. Ahmad, S., N. Singh and N.M. Syeda. 2020. Hydrochemical characteristics of the groundwater in Trans-Yamuna alluvial aquifer, Palwal district, Haryana. *Appl. Water Sci.* DOI:10.1007/s13201-020-1150-2.
21. Ministry of Water Resource. 1997. Ground resource estimation methodology. Report of the Groundwater Resource Estimation Committee, Government of India, New Delhi.
22. Qinghai, G., *et al.* 2007. A new model (DRARCH) for assessing groundwater vulnerability to arsenic contamination at basin scale : A case study in Taiyuan basin, Northern China. *Env. Geol.*, 52:923-932.
23. CGWB. 1982. Upper Yamuna project (1977-1982) report. Central Groundwater Board, North Western Region, Chandigarh.
24. CGWB. 2005. Groundwater exploration in Haryana State (as on 31st March 2004). Central Groundwater Board, North Western Region, Chandigarh.
25. Alam, F. 2010. Aquifer system and groundwater resource evaluation in parts of Hindon-Yamuna watershed in parts of western Uttar Pradesh. Ph.D. Thesis. Aligarh Muslim University, Aligarh.
26. Babiker, I.S., *et al.* 2005. A GIS based DRASTIC model for assessing aquifer vulnerability Kakamigahara Heights, Gifu Prefecture, Central Japan. *Sci. Total Env.*, 345 (1):127-140.
27. APHA. 1998. Standard methods for the examination of water and wastewater (20th edn). American Public Health Association, Washington D.C.

Development Of Process Parameters For The Clean Synthesis Of Silver Nanoparticles Using The Pernicious Aquatic Weed *Eichhornia crassipes* And An Assessment Of Their Properties

S. U. Ganaie¹, Tasneem Abbasi^{1*}, Tabassum Abbasi², R. Rajalakshmi¹ and S. A. Abbasi¹

1. Pondicherry University, Centre for Pollution Control and Environmental Engineering, Puducherry - 605 014, India

2. University of Petroleum and Energy Studies, Environmental Research Institute, Dehradun - 248 007, India

*Corresponding author, Email : abbasi.cpee@gmail.com

The effect of several process parameters on the biomimetic synthesis of silver nanoparticles (AgNPs) using ubiquitous and pernicious aquatic weed water hyacinth (*Eichhornia crassipes*) was studied. Aqueous extracts of the plant across a wide range of concentrations were able to successfully induce AgNP formation as well as their stabilization. It was shown with the aid of UV-visible spectroscopic, electron microscopic and dynamic light scattering studies that by manipulating the extract-Ag (I) stoichiometry, temperature, pH and interaction time, different shapes and sizes of nanoparticles can be generated. Conditions under which optimum formation of AgNPs of different shapes and sizes could be achieved were worked out. The efficacy of the AgNPs in free radical scavenging and catalysis was witnessed. The studies thus provide a template for scaling up the AgNP synthesis and utilizing the AgNPs. Given the fact that *E. crassipes* is freely available in large quantities, with no other recognized use, the present method opens up a possibility for large-scale utilization of it in synthesizing AgNPs of tunable shapes and sizes in a rapid, non-polluting, energy frugal and inexpensive manner.

KEYWORDS

Silver nanoparticles, Water hyacinth, *Eichhornia crassipes*, Biomimetic synthesis, Antioxidant activity, Catalysis

REFERENCES

1. Patil, M. P. and G. D. Kim. 2017. Eco-friendly approach for nanoparticles synthesis and mechanism behind antibacterial activity of silver and anticancer activity of gold nanoparticles. *Appl. Microbiol. Biotech.*, 101:79-92.
2. Singh, P., et al. 2016. Biological synthesis of nanoparticles from plants and microorganisms. *Trends Biotech.*, 34:588-599.
3. Fahmy, H. M., et al. 2017. Alternative green chemistry methods of silver nanoparticles synthesis : Review and comparison. *J. Bionanosci.*, 11:7-16.
4. Mondal, S., et al. 2014. A brief introduction to the development of biogenic synthesis of metal nanoparticles. *J. Nano Res.*, 27:41-52.
5. Shankar, S. S., et al. 2004. Rapid synthesis of Au, Ag and bimetallic Au core-Ag shell nanoparticles using neem (*Azadirachta indica*) leaf broth. *J. Colloid Interface Sci.*, 275:496-502.
6. Narayanan, K. B. and N. Sakthivel. 2011. Green synthesis of biogenic metal nanoparticles by terrestrial and aquatic phototrophic and heterotrophic eukaryotes and biocompatible agents. *Adv. Colloid Interface Sci.*, 169:59-79.
7. Gan, P. P. and S. F. Y. Li. 2012. Potential of plant as a biological factory to synthesize gold and silver nanoparticles and their applications. *Reviews Env. Sci. Biotech.*, 11:169-206.
8. Rao, Y. S., et al. 2013. Green synthesis and spectral characterization of silver nanoparticles from lakshmitulasi (*Ocimum sanctum*) leaf extract. *Spectrochimica Acta Part A: Molecular Biomo-lecular Spectroscopy*. 103: 156-159.

9. Abbasi, T., *et al.* 2015. Biomimetic synthesis of nanoparticles using aqueous extracts of plants (botanical species). *J. Nano Res.*, 31:138-202.
10. Chung, I. M., *et al.* 2016. Plant-mediated synthesis of silver nanoparticles: Their characteristic properties and therapeutic applications. *Nanoscale Res. Letters.* 11:1-14.
11. Abbasi, S. A. and P. C. Nipanay. 1986. Infestation by aquatic weeds of the fern genus *Salvinia* - Its status and control. *Env. Conser.*, 13:235-241.
12. Malik, A. 2007. Environmental challenge vis a vis opportunity: The case of water hyacinth. *Env. Int.*, 33:122-138.
13. Mishra, S. and A. Maiti. 2017. The efficiency of *Eichhornia crassipes* in the removal of organic and inorganic pollutants from wastewater: A review. *Env. Sci. Poll. Res.*, 24:7921-7937.
14. Guna, V., *et al.* 2017. Water hyacinth: A unique source for sustainable materials and products. *ACS Sustainable Chem. Eng.*, 5:4478-4490.
15. Mathur, S.M. 2007. Water hyacinth and its management through alternative utilization. ASABE - International Symposium on Air quality and waste management for agriculture.
16. Patel, S.A. 2011. Weed with multiple utility : *Lantana camara*. *Reviews Env. Sci. Biotech.*, 10:341–351.
17. Bortolotto, I. M. and N.G. Guarim. 2005. The use of the camalote, *Eichhornia crassipes* (Mart.) solms, Pontederiaceae, for handicraft in the district of Albuquerque, Corumba, MS, Brazil. *Acta Botanica Brasiliica.* 19:331–337.
18. Yan, S. H., *et al.* 2017. Advances in management and utilization of invasive water hyacinth (*Eichhornia crassipes*) in aquatic ecosystems- A review. *Critical Reviews Biotech.*, 37:1-11.
19. Rakotoarisoa, T., *et al.* 2016. Turning a problem into profit: Using water hyacinth (*Eichhornia crassipes*) for making handicrafts at Lake Alaotra, Madagascar. *Economic Botany.* 70:365-379.
20. Sekomo, C. B., *et al.* 2012. Heavy metal removal by combining anaerobic upflow packed bed reactors with water hyacinth ponds. *Env. Tech.*, 33(10-12):1455–1464.
21. Saravanan, S.P., *et al.* 2015. Textile wastewater treatment by phytoremediation efficiencies of water hyacinth (*Eichhornia crassipes*). *J. Chem. Pharmaceutical Sci.*, 8(4):790-792.
22. Sanmugapriya, E. and P. Senthamilselvan. 2017. Water hyacinth (*Eichhornia crassipes*) – An efficient and economic adsorbent for textile effluent treatment - A review. *Arabian J. Chem.*, 10: S3548-S3558.
23. Tripathi, B. and S. Shukla. 1991. Biological treatment of wastewater by selected aquatic plants. *Env. Poll.*, 69:69-78.
24. Tham, H. and P. Uden. 2013. Effect of water hyacinth (*Eichhornia crassipes*) silage on intake and nutrient digestibility in cattle fed rice straw and cottonseed cake. *Asian-Australasian J. Animal Sci.*, 26:646-653.
25. Thaeruzzaman, Q. and D. Kushari. 1989. Evaluation of some common aquatic macrophytes cultivated in enriched water as possible source of protein and biogas. *Hydrobiol. Bulletin.* 23:207-212.
26. Mako, A., *et al.* 2011. An evaluation of nutritive value of water hyacinth (*Eichhornia crassipes* Mart. solms-laubach) harvested from different water sources as animal feed. *Livestock Res. Rural Develop.*, 23:10.
27. Hontiveros, G. J. S. 2015. Nutritional value of water hyacinth (*Eichhornia*). *AAFL Bioflux.* 8:26–33.
28. Chaisaena, W., *et al.* 2012. Suitability of water hyacinth, rice straw and sunflower residues for the production of the edible mushrooms *Coprinopsis cinerea* and *Volvariella volvacea* in Thailand. *Chiang Mai University J. Natural Sci.*, 11(1):395-402.
29. Das, R. K., *et al.* 2016. Checking the biocom-patibility of plant-derived metallic nanoparticles: Molecular perspectives. *Trends Biotech.*, 34: 440-449.
30. Cheng, J., *et al.* 2010. Cogeneration of H₂ and CH₄ from water hyacinth by two-step anaerobic fermentation. *Int. J. Hydrogen Energy.* 35:3029–3035.
31. Chuang, Y. S., *et al.* 2011. Biohydrogen and biomethane from water hyacinth (*Eichhornia crassipes*) fermentation: Effects of substrate concentration and incubation temperature. *Int. J. Hydrogen Energy.* 36:14195–14203.
32. Abbasi, S. A., *et al.* 1990. Bioenergy potential of eight common aquatic weeds. *Biol. Wastes.* 34:359-366.

33. Ganesh, P. S., *et al.* 2005. Extraction of volatile fatty acids (VFAs) from water hyacinth using inexpensive contraptions and the use of the VFAs as feed supplement in conventional digesters. *Biochem. Eng. J.*, 27:17-23.
34. Tauseef, S. M., *et al.* 2013. Methane capture from livestock manure. *J. Env. Manage.*, 117:187-207.
35. Gajalakshmi, S., *et al.* 2001. Assessment of sustainable vermiconversion of water hyacinth at different reactor efficiencies employing *Eduriluseu geniae* Kinberg. *Bioresour. Tech.*, 80: 131-135.
36. Gajalakshmi, S., *et al.* 2002. High-rate composting- vermicomposting of water hyacinth. *Bioresour. Tech.*, 83: 235-239.
37. Abbasi, T. and S. A. Abbasi. 2010. Production of clean energy by anaerobic digestion of phytomass- New prospects for a global warming amelioration technology. *Renewable Sustainable Energy Reviews.* 141:653-1659.
38. Thombre, R., *et al.* 2014. A facile method for synthesis of biostabilized silver nanoparticles using *Eichhornia crassipes* (Mart.) Solms (water hyacinth). *Indian J. Biotech.*, 13:337-341.
39. Borase, P. H., *et al.* 2014. Plant extract: A promising biomatrix for eco-friendly, controlled synthesis of silver nanoparticles. *Appl. Biochem. Biotech.*, 173:1-29.
40. Ahmed, S., *et al.* 2016. Biosynthesis of gold nanoparticles: A green approach. *J. Phytochem. Photobiol., B: Biol.*, 161:141-153.
41. Anuradha, J., *et al.* 2015. An eco-friendly method of synthesizing gold nanoparticles using an otherwise worthless weed pistia (*Pistia stratiotes* L.). *J. Adv. Res.*, 6:711–720.
42. Ganaie, S. U., *et al.* 2014. Biomimetic synthesis of silver nanoparticles using the amphibious weed ipomoea and their application in pollution control, *J. King Saud University - Sci.*, 26:222-229.
43. Ganaie, S. U., *et al.* 2015. Green synthesis of silver nanoparticles using an otherwise worthless weed mimosa (*Mimosa pudica*): Feasibility and process development toward shape/size control. *Particulate Sci. Tech.*, 33:638-644.
44. Ganaie, S. U., *et al.* 2016. Low-cost, environment-friendly synthesis of palladium nanoparticles by utilizing a terrestrial weed *Antigonon leptopus*. *Particulate Sci. Tech.*, 34:201–208.
45. Mashwani, Z. R., *et al.* 2016. Applications of plant terpenoids in the synthesis of colloidal silver nanoparticles. *Adv. Colloid Interface Sci.*, 234: 132-141.
46. Sreelatha, S. and P. R. Padma. 2009. Antioxidant activity and total phenolic content of *Moringa oleifera* leaves in two stages of maturity. *Plant Foods Human Nutrition.* 64(4):303-311.
47. Abdel-Aziz, M. S., *et al.* 2014. Antioxidant and antibacterial activity of silver nanoparticles biosynthesized using *Chenopodium murale* leaf extract. *J. Saudi Chem. Society.* 18(4):356-363.
48. Muniyappan, N. and N.S. Nagarajan. 2014. Green synthesis of silver nanoparticles with *Dalbergia spinosa* leaves and their applications in biological and catalytic activities. *Process Biochem.*, 49: 1054-1061.

Identification Of Groundwater Prospective Zones Of Varuna Watershed Using Weighted Overlay Method

Vikram Kumar^{1*}, Kailash Narayan², Manvendra Singh Chauhan³, Prabhat K. S. Dikshit⁴ and Shyam B. Dwivedi⁴

1. Gaya College of Engineering, Department of Civil Engineering, Gaya, Bihar, India

2. Delhi Technical Campus, Department of Civil Engineering, Greater Noida, India

3. Holy Mary Institute of Technology and Science, Department of Civil Engineering, Hyderabad, India

4. Indian Institute of Technology (BHU), Department of Civil Engineering, Varanasi – 221 005, U.P., India

In this work, potential groundwater zones of the Varuna watershed have been identified using the Weighted Index Overlay method. To apply this method, various thematic layers, soil, drainage, slope, land use land cover, and topographic layers have been considered. The comparative weight has been assigned to individual thematic layers and further rank assigned to every category of thematic layers. The overlying of layers has been done in ArcGIS to produce a potential groundwater zones map. Potential groundwater zones (PGZs) of the Varuna watershed have been categorized into three categories: good, moderate and low. Results show that the maximum part of the study area has a moderate groundwater zone. The criterion to categorize PGZs is based on the depth of the groundwater table from the ground surface. This study shows that remote sensing and GIS are the most useful tools to explore the groundwater potential zones and opened new paths to take care of the water resources.

KEYWORDS

Varuna watershed, GIS, Potential groundwater zones

REFERENCES

1. Narendra, K., K. Nageswararao and P. Swarnalatha. 2013. Integrating remote sensing and GIS for identification of groundwater prospective zone in the Narava basin Visakhapatnam region, Andhra Pradesh. *J. Geol. Society India*. 81:248-260.
2. Chennai, I. and S. Khemiri. 2009. Evaluation of groundwater quality using multiple linear regression and structural equation modeling. *Int. J. Env. Sci. Tech.*, 6:509-519.
3. Kumar, V. and S. Sen. 2018. Evaluation of spring discharge dynamics using recession curve analysis : A case study in data-scarce region, Lesser Himalayas, India. *Sustainable Water Resour. Manage.*, 4(3):539-557.
4. Kumar, V. and S. Sen. 2020. Assessment of spring potential for sustainable agriculture : A case study in Lesser Himalayas. *Appl. Eng. Agric.*, 36(1):11-24.
5. Kumar, V. and S. Paramanik. 2020. Application of high-frequency spring discharge data : A case study of Mathamali spring rejuvenation in Garhwal Himalaya. *Water Supply*.
6. Apel, M. 2006. From 3D geomodeling systems towards 3D geoscience information systems : Data model query functionality and data management. *Computers Geosci.*, 32(2):222-229.
7. Yoo, C. and J.M. Kim. 2007. Tunneling performance prediction using integrated GIS and neural network. *Computers Geotechnics*. 34(1):19-30.
8. Sreedhar, G., et al. 2009. Mapping of groundwater potential zones in the Musi basin using remote sensing data and GIS. *Adv. Eng. Software*. 40:506-518.
9. Chowdhury, A., M.K. Jha and V.M. Chowdhury. 2010. Delineation of groundwater recharge zones and identification of artificial recharge sites in west Medinipur district, West Bengal using RS GIS and MCDM techniques. *Env. Earth Sci.*, 59 (6):1209-1222.
10. Malik, M. I., M. S. Bhat and S. A. Najor. 2016. Remote sensing and GIS-based groundwater potential mapping for sustainable water resource management of ladder catchment in Kashmir valley. *J. Geol. Society India*. 87(6):631-760.

11. Siva, G., N. Nasir and R. Selvakumar. 2017. Delineation of groundwater potential zone in Sengipatti for Thanjavur district using analytical hierarchy process. *IOP Conference Series : Earth and Env. Sci. Earth.* 80(1):2063.
12. Nag, S.K. and A. Kundu. 2018. Application of remote sensing, GIS and MCA techniques for delineating groundwater prospect zones in Kashipur block, Purulia district, West Bengal. *Appl. Water Sci.*, 8.
13. Raviraj, A., N. Kuruppath and B. Kannan. 2017. Identification of potential groundwater recharge zones using remote sensing and geographical information system in Amaravathy basin. *J. Remote Sensing GIS.* 6(4):213.
14. Tabesh, M., A.H.A. Yekta and R. Burrows. 2009. An integrated model to evaluate losses in water distribution systems. *Water Resour. Manage.*, 23:477-492.
15. Choudhury, V.M., H.H. Rao and P.B.S. Sharma. 2015. GIS-based decision support system for groundwater assessment in large irrigation project areas. *Agric. Water Manage.*, 3:229-252.
16. Adham, M.I., et al. 2010. Study on groundwater recharge potentiality of baring tract Rajshahi district, Bangladesh using GIS and remote sensing technique. *J. Geol. Society India.* 7:432-438.
17. Gupta, M. and P.K. Srivastava. 2010. Integrating GIS and remote sensing for identification of groundwater potential zones in the hilly terrain of Pavagarh, Gujarat. *Water Int.*, 35:233-245.
18. Vijith, H., et al. 2012. An assessment of soil erosion probability and erosion rate in a tropical mountainous watershed using remote sensing and GIS. *Arabian J. Geosci.*, 5:797-805.
19. Subla, R.N. 2009. A numerical scheme for groundwater development in a watershed basin of basement terrain : A case study from India. *Hydrogeol. J.*, 17:379-396.
20. Olutoyin, A., et al. 2014. Delineation of groundwater potential zones in the crystalline basement terrian of SW-Nigeria : An integrated GIS and remote sensing approach. *Appl. Water Sci.*, 4(1):19-38.
21. Singh, S.M. 2014. Morphology changes of Ganga river over time at Varanasi. *J. River Eng.*, 2:2.
22. Jaiswal, J. and N. Verma. 2013. The study of landuse/land cover in Varanasi district using remote sensing and GIS. *Transaction Institute Indian Geographers.* 35(20):201-208.
23. Narayan, K., P.K.S Dikshit and S.B. Dwivedi. 2012. GIS supported geomorphologic instantaneous unit hydrograph (GIUH) of Varuna river basin using geomorphological characteristics. *Int. J. Adv. Earth Sci.*, 1(2):68-76.
24. Lillesand, T.M. and R.W. Kiefer. 2002. Remote sensing and image interpretation. John Wiley and Sons (Asia) Pvt. Ltd., Singapore.
25. Kumar and Shankar. 2014. Assessment of groundwater potential zones using GIS. *Frontiers Geosci.*, 2(1):1-10.
26. Preeja, K.R., et al. 2011. Identification of groundwater potential zones of a tropical river basin (Kerala) using remote sensing and GIS techniques. *J. Indian Society Remote Sensing.* 39:83-94.
27. Edet, A.E., et al. 1998. Application of remote sensing data groundwater exploration : A case study of the cross-river state southeastern Nigeria. *Hydrogeol. J.*, 6(3):394-404.
28. Magesh, N.S. and N. J. Chandrasekar. 2012. Delineation of groundwater potential zones in Theni district, Tamil Nadu using remote sensing, GIS and MIF techniques. *Geosci. Front.*, 3(2):189-196.
29. Waikar, M.L. and A.P. Nilawar. 2014. Identification of groundwater potential zone using remote sensing and GIS techniques. *Int. J. Innovative Res. Sci. Eng. Tech.*, 3(5):12163-12174.
30. Horton, R.E. 1945. Erosional development of streams and their drainage basins : Hydrophysical approach to quantitative morphology. *Geol. Society America Bulletin.* 56:275-370.
31. Wisler, C.O. and B.F. Brater. 1959. Hydrology. New York.
32. Willey. D.K. and M.L.W. Todel. 2005. Groundwater hydrology. John Wiley and Sons, Hoboken.
33. Pani, S., A. Chakraborty and S. Bhadury. 2016. Groundwater potential zone identification by analytical hierarchy process (AHP) weighted overlay in GIS environment - A case study of Jhargam block, Paschim Medinipur. *Int. J. Remote Sensing Geosci.*, 5(3):1-10.
34. Zeyad, M. 2011. Using remote sensing approach and surface landscape conditions for optimization of watershed management in Mediterranean regions. *Phys. Chem. Earth.* 36:213-220.
35. Krishnamurthy, J., et al. 1996. An approach to demarcate groundwater potential zones through remote sensing and a geographical information system. *Int. J. Remote Sensing.* 17(10):1867-1884.
36. Prasad, R.K., et al. 2008. Deciphering potential groundwater zone in hard rock through the application of GIS. *Env. Geol.*, 55:467-475.
37. Muheeb, M.A. and A.J. Rasheed. 2009. Evaluation of aquifers vulnerability to contamination in the Yarmouk river watershed, Jordan, based on drastic method. *Arabian J. Geosci.*, 3:273-282.

Effect Of Superplasticizer On Alkali Activated Ground Granulated Blast Furnace Slag Concrete In Ambient Curing Condition For Sustainable Environment

K. Naga Rajesh¹, P. Markandeya Raju^{2*} and Kapileswar Mishra¹

1. Centurion University of Technology and Management, Department of Civil Engineering, Bhubaneswar, Odisha, India

2. MVGR College of Engineering (A), Department of Civil Engineering, Vizianagaram, India

*Corresponding author, Email : markandeyaraju@yahoo.com; rajeshkanta@gmail.com

Concrete is the most widely used construction material due to its mechanical and durability properties. Due to the use of ordinary portland cement in concrete production, CO₂ emissions occur from cement leading to environmental pollution. The objective of the current study is to minimize the cement content in the concrete production. In this regard, cement is replaced with ground granulated blast furnace slag, alkaline solution is used for alkali activation to develop geopolymerization process and this type of concrete produced is called as alkali-activated slag based geopolymer concrete. Most of the research works available are related to heat-cured flyash based geopolymer concrete and less work on ambient cured. The present work is one such alternative method in producing ambient cured ground granulated blast furnace slag geopolymer concrete. The compressive strength of ambient cured slag based geopolymer concrete with and without superplasticizer is 5.5% and 8.7% higher than ordinary portland cement concrete, respectively.

KEYWORDS

Geopolymer concrete, Ambient cured geopolymer concrete, Slag based geopolymer concrete, Superplasticizer geopolymer concrete, Sustainable concrete

REFERENCES

1. EPA. 2010. Available and emerging technologies for reducing greenhouse gas emissions from the portland cement industry. Environmental Protection Agency, Washington, D.C.
2. Van Oss, H.G. 2005. Background facts and issues concerning cement and cement data. U.S. Geological Survey, Reston, VA.
3. Vishwakarama, V. and D. Ramachandran. 2018. Green concrete mix using solid waste and nanoparticles as alternative - A review. *Construction Building Mater.*, 162:96-103.
4. Liew, K.M., A.D. Sojobi and L.W. Zhang. 2017. Green concrete : Prospects and challenges. *Construction Building Mater.*, 156:1063-1095.
5. Davidovits, J. 2013. Geopolymer cement : A review. *Geopolymer Sci. Technics.*
6. Murtazaev, S.A.Y., et al. 2018. The use of highly active additives for the production of clinkerless binders. International Symposium on Engineering and earth science : Applied and fundamental research (ISEES 2018). DOI : 10.2991/isees-18.2018.68.
7. Shi, C., A.F. Jimenez and A. Palomo. 2011. New cements for the 21st century : The pursuit of an alternative to portland cement. *Cement Concrete Res.*, 41 (7):750-763. DOI:10.1016/j.cemconres.2011.03.0.16.
8. IS 455. 1989. Portland slag cement-Specification (reaffirmed 1995). Bureau of Indian Standard, New Delhi.
9. IS 10289. 1987. Specification for granulated slag for the manufacture of portland slag cement (reaffirmed 1999). Bureau of Indian Standards, New Delhi.
10. IS 2720. 1980. Part 3: Methods of test for aggregates for specification (reaffirmed 2002). Bureau of Indian Standards, New Delhi.

11. IS 2386. 1963. Methods of test of aggregates for concrete : Part 3. Specific gravity, density, voids, absorption and bulking. Bureau of Indian Standards, New Delhi.
12. IS 2386. 1963. Methods of test for aggregates for concrete : Part 1. Particle size and shape. Bureau of Indian Standards, New Delhi.
13. IS 2645. 2003. Integral water proofing compounds for cement mortar and concrete-Specification (reaffirmed 2005). Bureau of Indian Standards, New Delhi.
14. Patankar, S.V., Y.M. Ghugal and S.S. Jamkar. 2014. Mix design of flyash based geopolymer concrete. *Adv. Structural Eng.*, 1619-1634. DOI:10.1007/978-81-322-2187-6_123.
15. Davidovits, J. Chemistry of geopolymeric systems, technologies. Proceedings of Second International Conference on geopolymer. Geopolymer Institute. pp 9-40.
16. Davidovits, J. 2008. Geopolymer chemistry and applications (2nd edn). Institute Ge'opolyme're, France.
17. Nath, P. and P.K. Sarker. 2014. Effect of GG BFS on setting, workability and early strength properties of flyash geopolymer concrete cured in ambient condition. *Construction Building Mater.*, 66:163-171. DOI:10.1016/j.conbuildmat. 2014. 05.080.
18. Nematollahi, B. and J. Saniyan. 2014. Effect of different superplasticizers and activator combinations on workability and strength of flyash based geopolymer. *Elsevier (Materials Design)*. 5(57): 667-672.
19. IS 456. 2000. Plain and reinforced concrete code of practice. Bureau of Indian Standards, New Delhi.
20. Nath, P. and P.K. Sarker. 2017. Flexural strength and elastic modulus of ambient-cured blended low-calcium flyash geopolymer concrete. *Construction Building Mater.*, 130:22-31. DOI:10.1016. j.conbuildmat. 2016.11.034.

Evolution Of Physico-Chemical Parameters And Pathogenic Microorganisms Of Sludge Of Dar El Gueddari Wastewater Treatment Stations (Morocco)

H. Griou*, S. Njimat, M. Aboulouafa and S. Ibn Ahmed

University Ibn Tofail, Laboratory of Materials, Electrochemistry and Environment, Faculty of Sciences, Department of Chemistry, Kenitra, Morocco

*Corresponding author, Email : hajarsmp@gmail.com

The purpose of this study is to evaluate the effect of the storage of sewage sludge in Geotube® on the physico-chemical and microbiological parameters of the sludge resulting from the cleaning of the anaerobic basins of the Dar El Gueddari treatment plant between October 2013 and May 2017. The results obtained during this study revealed an increase in the pH of sludge from 6.5-7.2 and a gradual reduction in odour according to the drying time. The dryness reaches an optimal value of 76%, the variation of the C/N ratio goes from 9.62-11.80 (lower than 15), which indicates that these sludges have fast mineralization of organic matter. For nutrients, there was a slight decrease except for total nitrogen, which dropped from 1.3 to 0.8 ppm. The contents of metallic trace elements (Cd, Cr, Cu, Hg, Pd and Zn) and the organic trace components [7 PCBs (28, 52, 101, 118, 138, 153, 180)], fluoranthene, benzo(a)fluoranthene, benzo(b)fluoranthene are in small quantities and well below the French regulatory standards. For the microbiological component, the obtained results showed a decrease in the bacterial load over time and it is eliminated completely from the second year. The so-called dewatering treatments applied to sludge play a major role in the elimination of most or all pathogenic micro-organisms.

KEYWORDS

Sewage sludge, Physico-chemical, Microbiological parameters, Wastewater

REFERENCES

1. Hiroux, G., G. Lefevre and M.A. Gaffet. 1983. Organic composition and decomposition in the soil of mixed sludge from an urban wastewater treatment plant. *Soil Sci.*, 1:17-26.
2. Krogmann, U. 2001. Composting : Sludge into biosolid : Processing, disposal and utilization. In Sous la direction. Ed L. Spinosa and P.A. Vesilind. Iwa Publishing, Londres, R.U.
3. Jebari, H. 2014. Sludge management from the Grand Nador wastewater treatment plant. Workshop on the management and recovery of sludge from ONEE-Water branch, Rabat, Morocco. Proceedings, pp 15.
4. Boudreault, P. and J.F. Blais. 2002. Optimization of STABIOX technology for the stabilization, deodorization and improvement of the dewatering of paper mill sludge. Workshop on R and D in waste recovery, Sainte-Foy Quebec. Proceedings, pp 122.
5. Haut-commissariat plan. 2014. General population census : City of Dar El Gueddari.
6. BRL. 2015. Update study of the master plan for integrated water resources development in Gharb plan. BRL ingeniere.
7. ISO 10390. 2005. Soil quality determination of pH.
8. AFNOR, NF ISO 11265, X 31-113. 1995. Normadoc.
9. BS EN 13342. 2000. Characterization of sludges. Determination of Kjeldahl nitrogen.
10. ISO 16649-2. 2000. Microbiology of food and animal feeding stuffs-Horizontal method for the enumeration of beta-glucuronidase-positive *Escherichia coli*-Part 2: Colony count technique at 44°C using 5-bromo-4-chloro-3-indolyl beta-D-glucuronide.
11. IMANOR NM ISO 6888-2. 2016. Food microbiology : Horizontal method for the enumeration of coagulase positive staphylococci *Staphylococcus aureus* and other species. Part 2 : Technique using gelose medium with rabbit plasma and fibrinogen.
12. ISO 11290-1. 2017. Microbiology of the food chain-Horizontal method for the detection and enumeration of *Listeria monocytogenes* and of *Listeria* spp. Part I : Detection method.

13. ISO/TS6579-2. 2012. Microbiology of food and animal feed–Horizontal method for the detection, enumeration and serotyping of Salmonella. Part 2: Enumeration by a miniaturized most probable number technique.
14. ONEE, 2013. Sludge spreading plan from the Dar El Gueddari WWTP.
15. Noble, C. 1997. Treat and recover sludge. OTV.
16. Rigby, H., *et al.* 2016. A critical review of nitrogen mineralization in biosolids-amended soil, the associated fertilizer value for crop production and potential for emissions to the environment. *Sci. Total Env.*, 541:1310-1338.
17. Tremblay, J., *et al.* 2008. Storage and long-term stability of municipal sewage sludge decontaminated and stabilized by chemical or biological means. *J. Env. Eng. Sci.*, 7(4):357-368.
18. Tu, J., *et al.* 2012. Heavy metal concentration and speciation of seven representative municipal sludges from wastewater treatment plants in northeast China. *Env. Monit. Assess.*, 184(3):1645-1655.
19. ONEE. 2004. Characterization study of wastewater in the city of Dar El Gueddari.
20. Berthou, M. 2010. Evaluation of the efficiency of filter gardens® for the treatment of urban and industrial sludge. Ph.D. Thesis. Universite 'Henri Poincare'; INPL-Institute National Polytechnique de Lorraine.

Total Suspended Particulate Matter And PM₁₀ Concentrations Related Meteorological Conditions In Daya, Makassar

Y. Sattar^{1*}, A. Fitri², A. Nani³, M. Ramdiana⁴ and A. Syarifuddin⁵

1. Universitas Muslim Indonesia, Department of Mechanical Engineering, Makassar - 90231, Indonesia

2. Universitas Bosowa, Department of Chemical Engineering, Makassar - 90231, Indonesia

3. Universitas Bosowa, Department of Environmental Engineering, Makassar - 90231, Indonesia

4. Universitas Muhammadiyah, Department of Urban and Regional Planning, Pare-Pare - 91131, Indonesia

5. Universitas Muslim Indonesia, Department of Electrical Engineering, Makassar - 90231, Indonesia

*Corresponding author, Email : sattar.teknik@umi.ac.id

Ambient total suspended particulate matter (TSP) and PM₁₀ (that is particulate diameter less than 10µm in size) produced by human activities, such as motorized vehicle emissions and industries can affect ambient air quality. On the other hand, the Makassar City Power area as a sampling site which is now turning into a densely populated area due to the rapid development of residential areas and this fact enables more opportunities to many residents and disruption of human health because of the decreased ambient air quality especially due to the presence of particulate matter, while the concentration of TSP and PM₁₀ is influenced by meteorological conditions. In this study, Pearson's coefficient of correlation was applied to study the relationship between TSP, PM₁₀ and meteorological variables, that is humidity, temperature, wind speed and rainfall. TSP and PM₁₀ sampling was done using the high volume air sampler (HVAS) tool, for meteorological factors using the hygrometer, thermometer and using anemometer, while rainfall data was obtained from the Office of Meteorological and Geophysics area IV Makassar. This study concluded that the temperature was found as a significant factor compared with other factors that influence the concentration of TSP and PM₁₀. Increased rainfall, humidity and wind speed have a negative correlation with the average concentration of TSP and PM₁₀ in Daya, Makassar.

KEYWORDS

Air pollutants, Particulate matter, Meteorological parameters, Statistical analysis, Makassar

REFERENCES

1. Rashid, M., *et al.* 2014. PM₁₀ black carbon and ionic species concentration of urban atmosphere in Makassar of South Sulawesi Province, Indonesia. *Atmos. Poll. Res.*, 5: 610-615. DOI: 10.5094/APR.2014.070.
2. Ganguly, R. and P. Kumar. 2018. The air quality assessment of northern hilly city in India. *Indian J. Env. Prot.*, 38 (12): 983-997.
3. Sharma, D. and U.C. Kulshresta. 2014. Spatial and temporal patterns of air pollutants in rural and urban area of India. *Env. Poll.*, 195: 276-281.
4. Pratt, G.C., *et al.* 2018. Measurements of gas and particle polycyclic aromatic hydrocarbons (PAHs) in air at urban, rural and near-roadway sites. *Atmos. Env.*, 179:268-278.
5. Kopanakis, I., *et al.* 2018. Impact from local sources and variability of fine particle number concentration in a coastal sub-urban site. *Atmos. Res.*, 213: 136-148.
6. Ayala, A., *et al.* 2012. Air pollutants and sources associated with health effects. *Air Quality Atmos. Health*. 5:151-167.
7. Dash, S.K., *et al.* 2019. Atmospheric dispersion modeling using aermოდ to predict the impact of PM₁₀ near Bileipada, Odisha. *Indian J. Env. Prot.*, 39(4): 299-309.
8. Rodrigues, F.A. and I. Joekes. 2011. Cement industry: Sustainability, challenges and perspective. *Env. Chem. Letter*. 9: 151-166.

9. Allaban, M.A. and H.A. Qudais. 2011. Impact assessment of ambient air quality by cement industry: A case study in Jordan. *Aerosol Air Quality Res.*, 11(7): 802-810.
10. Hassan, S. K. 2018. Sources and cancer risk of heavy metal in total suspended particulate in some square areas of Greater Cairo, Egypt. *Indian J. Env. Prot.*, 38(12): 1040-1050.
11. Hoek, G., *et al.* 2013. Long-term air pollution exposure and cardio-respiratory mortality: A review. *Env. Health*. DOI:10.1186/1476-069X-12-43.
12. Maslh, A. 2020. Additive regression algorithm predicts the atmospheric pollutant concentration with higher precision. *Indian J. Env. Prot.*, 40(3): 253-258.
13. Sattar, Y., *et al.* 2019. Characteristics of the PM₁₀ in the urban environment of Makassar, Indonesia. *J. Urban Env. Eng.*, 13(1): 198-207. DOI: 10.4090/juee.2009.v13n1.198-207.
14. Sattar, Y., *et al.* 2014. Black carbon and elemental concentration of ambient particulate matter in Makassar, Indonesia. *IOP Conf. Series: Earth Env. Sci.*, 18. DOI:10.1088/1755-1315/18/1/012099.
15. Stafoggia, M., *et al.* 2016. Particle number concentrations near the Rome-Ciampino city airport. *Atmos. Env.*, 147: 264-273.
16. Masiol, M., *et al.* 2017. Analysis of major air pollutants and submicron particles in New York city and Long Island. *Atmos. Env.*, 148: 203-214.
17. Saini, M., *et al.* 2018. The influence of throat length and vacuum pressure on air pollutant filtration using ejector. AIP Conference Proceedings 1977. DOI: 10.1063/1.5042939.
18. Grivas, G. and A. Chaloulakou. 2006. Artificial neural network models for prediction of PM₁₀ hourly concentrations, in the Greater area of Athens, Greece. *Atmos. Env.*, 40:1216-1229.
19. Carnevale, C., *et al.* 2010 A non-linear analysis to detect the origin of PM₁₀ concentrations in Northern Italy. *Sci. Total Env.*, 409: 182-191.
20. Giri, D., *et al.* 2008. The influence of meteorological conditions on PM₁₀ concentrations in Kathmandu valley. *Int. J. Env. Res.*, 2: 49-60.
21. EPA. 2009. Assessment of the impacts of global change on regional U.S. air quality: A synthesis of climate change impacts on ground-level ozone. Environmental Protection Agency, Washington, DC.
22. Kastner, K.P. and M.W. Rotach. 2004. Mean and monthly turbulence characteristics in an urban roughness sub-layer. *Boundary Layer Meteorol.*, 111: 55-84.
23. Jacob, D.J. and D.A. Winner. 2009. Effect of climate change on air quality. *Atmos. Env.*, 43:51- 63.
24. Karppinen, A., S.M. Joffre and J. Kukkonen. 2000. The refinement of a meteorological pre-processor for the urban environment. *Int. J. Env. Poll.*, 14: 565-572.
25. Roy, A., *et al.* 2019. Particulate matter and its influence on air quality for Mumbai and Agra. *Indian J. Env. Prot.*, 39(12): 1075-1083.
26. DOE. 2002. Environment status report. Department of Environment, Indonesia.
27. DOE. 1999. Air pollution control. Government decree no. 41/1999. Department of Environment, Indonesia.
28. Faizal, F. and A.M. Ulfah. 2009. Correlation between total rainfall and TSP levels in 2004-2008 in Jakarta in the process of cleaning the atmosphere by rainfall. *Bulletin Meteorol. Climatol. Geophysics*. 5(3): 263-274.
29. Mkoma, S.L. and I.C. Mjemah. 2011. Influence of meteorology on ambient air quality in Morogoro, Tanzania. *Int. J. Env. Sci.*, 1(6): 1107-1115.
30. Sattar, A., *et al.* 2012. A preliminary survey of air quality in Makassar city South Sulawesi, Indonesia. *J. Teknologi* (1). 57.
31. Rahman, S.A., *et al.* 2011. Sources apportionment of fine and coarse aerosol in Klang valley, Kualalumpur using positive matrix factorization. *Atmos. Poll. Res.*, 2: 197-206.
32. Kothai, P., *et al.* 2008. Source apportionment of coarse and fine paarticulate matter at Navi Mumbai, India. *Aerosol Air Quality Res.*, 70: 426-436.
33. Santoso, M., *et al.* 2008. Source identification of the atmospheric aerosol at urban and suburban in Indonesia by positive matrix factorization. *Sci. Total Env.*, 397: 229-237.
34. Gvodic', V., *et al.* 2011. Influence of meteorological factors NO₂, SO₂, CO and PM₁₀ on the concentration of O₃ in the urban atmosphere of Eastern Croatia. *Env. Model Assess.*, 16: 491-501.
35. Akpınar, S., *et al.* 2008. Evaluation of relationship between meteorological parameters and air pollutant concentrations during winter season in Elazyđ, Turkey. *Env. Monit. Assess.*, 146: 211-224.

Identification And Characterization Of Microplastics At Muara Gembong, Bekasi District, Indonesia

Uswah Hasanah^{1*}, Haeruddin² and Pujiono Wahyu Purnomo²

1. Diponegoro University, Master of Coastal Resources Management, Semarang, Indonesia

2. Diponegoro University, Department of Aquatic Resources, Semarang, Indonesia

*Corresponding author, Email : uswahtr@gmail.com

One of the factors causing sea pollution is a large amount of plastic waste. Over the past 40 years, the world's plastic production has increased by 25 times and the material recovered is no more than 5%. According to the World Bank in 2018, Indonesia is a contributor to 9 million tonnes of plastic waste annually. Plastic waste in the sea will be cut into smaller sizes or called microplastics, over 100 years. Microplastics have an impact that can save the lives of marine life. Globally, the growth of plastics has increased significantly, in the 1950s the amount of plastic reached 1.7 million tonnes and in 2014 it had increased to 299 tonnes. Muara Gembong is part of the downstream of the Citarum river in Bekasi Regency which has a dense population and has the potential to contribute to plastic waste in the sea. Based on the results of research, there are four types of microplastics found, namely fibres fragments, films and pellets. Microplastic characterizations found were polypropylene (PP), polystyrene (PS), nitrile, nylon, low-density polyethylene (LDPE) and linear low-density polyethylene (LLDPE) and high-density polyethylene (HDPE). This can happen because of the dense population which will increase the activities of the community around the coastal area.

KEYWORDS

Microplastic, FTIR, Muara Gembong

REFERENCES

1. Dewi, S. I., A. A. Budiarsa and I. R. Ritonga. 2015. Microplastic distribution of sediments in Muara Badak, Kutai Kartanegara Regency. *Depik.*, 4(3): 121–131.
2. Sutherland, W. J., *et al.* 2010. A horizon scan of global conservation issues for 2010. *Trends Ecol. Evol.*, 25(1): 1-7.
3. Kershaw, P. J. 2015. Biodegradable plastics and marine litter : Misconceptions, concerns and impacts on marine environments. United Nations Environment Programme.
4. Anonymous. 2015. Plastics-The facts : An analysis of European plastics production, demand and waste data. PlasticsEurope, Brussels-Belgium.
5. Khoironi, A., S. Anggoro and Sudarno. 2018. The existence of microplastic in Asian green mussels. *IOP Conf. Ser.: Earth Env. Sci.*, 131(1).
6. Cordova, M. R., A. I. S. Purwiyanto and Y. Suteja. 2019. Abundance and characteristics of microplastics in the northern coastal waters of Surabaya, Indonesia. *Mar. Poll. Bull.*, 142(3): 183-188.
7. National Oceanic and Atmospheric Administration Marine Debris Programme. 2016. Report on marine debris impacts on coastal and benthic habitats. NOAA Marine Debris Programme, Silver Spring, Maryland.
8. Wright, S. L., R. C. Thompson and T. S. Galloway. 2013. The physical impacts of microplastics on marine organisms: A review. *Env. Poll.*, 178: 483-492.
9. Browne, M. A., *et al.* 2013. Microplastic moves pollutants and additives to worms, reducing functions linked to health and biodiversity. *Curr. Biol.*, 23(23): 2388-2392.
10. Hiwari, H., *et al.* 2019. Condition of microplastic garbage in sea surface water at around Kupang and Rote, East Nusa Tenggara Province. Proceedings of National Seminar on Indonesian biodiversity society. 5:165-171.
11. Ayuningtyas, W. C. 2019. Microplastic abundance in waters in Banyuurip, Gresik, East Java. *JFMR-J. Fish. Mar. Res.*, 3 (1):41-45.
12. Cordova, M. R. and A. J. Wahyudi. 2016. Microplastic in the deep-sea sediment of southwestern Sumatran waters. *Marine Res. Indonesia*. 41(1): 27-36.

13. Van Cauwenberghe, L., *et al.* 2013. Microplastic pollution in deep-sea sediments. *Env. Poll.*, 182: 495–499.
14. Tagg, A. S., *et al.* 2015. Identification and quantification of microplastics in wastewater using focal plane array-based reflectance micro-FTIR imaging. *Anal. Chem.*, 87(12): 6032-6040.
15. Nor, M. H. and J. P. Obbard. 2014. Microplastics in Singapore's coastal mangrove ecosystems. *Mar. Poll. Bull.*, 79(2):278-283.
16. Claessens, M., *et al.* 2011. Occurrence and distribution of microplastics in marine sediments along the Belgian coast. *Mar. Poll. Bull.*, 62(10):2199-2204.
17. Hafidh, D., I. W. Restu and N. Made. 2018. Study of microplastic abundance in Bena Bay waters of Bali Province. *Curr. Trends Aquat. Sci.*, 88: 80-88.
18. Hastuti, A. R., F. Yulianda and Y. Wardianto. 2014. Spatial distribution of marine debris in mangrove ecosystem of Pantai Indah Kapuk, Jakarta. *Bonorowo Wetl.*, 4(2): 94-107.
19. Deepika, S. and R. J. Madhuri. 2015. Biodegradation of low density polyethylene by micro-organisms from garbage soil. *J. Exp. Biol. Agric. Sci.*, 3(1): 15-21.
20. Jung, M. R., *et al.* 2018. Validation of ATR FTIR to identify polymers of plastic marine debris, including those ingested by marine organisms. *Mar. Poll. Bull.*, 127(11):704-716.
21. Andrady, A. L. 2011. Microplastics in the marine environment. *Mar. Poll. Bull.*, 62(8): 1596-1605.
22. Jambeck, J. R., *et al.* 2015. Plastic waste inputs from land into the ocean. *Sci.*, 347(6223):768-771.
23. Caruso, G. 2015. Plastic degrading microorganisms as a tool for bioremediation of plastic contamination in aquatic environments. *J. Poll. Eff. Cont.*, 3(3):9-11.
24. Asmita, K., T. S. Singh and S. Tejashree. 2015. Isolation of plastic degrading micro-organisms from soil samples collected at various locations in Mumbai, India. *Int. Res. J. Env. Sci.*, 4(3): 77-85.
25. Trivedi, P., *et al.* 2016. Role of microbes in degradation of synthetic plastics and manufacture of bioplastics. *J. Chem. Pharm. Res.*, 8(3):211-216.

Anticipated Performance Index Of Tree Species As An Indicator For Green Belt Development In Traffic Density Area

Gopamma D.*, Jagadeeswara Rao K., Suresh Kumar K. and Srinivas N.

GITAM (Deemed to be University), Department of Environmental Science, Institute of Science, Visakhapatnam, Andhra Pradesh, India

* Corresponding author, Email: gdaka@gitam.edu

Air pollution is one of the major environmental problems in the urban atmosphere and green plants act as living filters to absorb and accumulate pollutants from the atmosphere. To assess the plants' tolerance, the parameters of air pollution tolerance index (APTI) and anticipated performance index (API) in the traffic density area of Visakhapatnam were evaluated. Seventeen tree species were identified growing in the selected study area. The APTI was calculated based on the biochemical analysis of leaf cell sap pH, total chlorophyll content, relative water content (RWC) and ascorbic acid. The air quality in traffic density areas ranged from moderate to high for particulate matter and low to moderate for NO_x. Out of 17 species studied, eight tree species (*Ficus benghalensis*, *Eucalyptus citriodora*, *Mangifera indica*, *Artocarpus heterophyllus*, *Syzygium cumini*, *Azadirachta indica*, *Bauhinia purpurea* and *Pongamia pinnata*) have shown API values above 81 and are categorized as excellent and can be grown in urban areas. Among them, *Azadirachta indica* and *Pongamia pinnata* are suitable for avenue plantation in traffic density areas because of their resistance to pollution and extreme winds during cyclones.

KEYWORDS

Urban plantation, Green belt, Anticipated performance index, Traffic density area

REFERENCES

- Jahan, S. and M. Zafar Iqbal. 1992. Morphological and anatomical studies on leaves of different plants affected by motor vehicle exhaust. *J. Islamic Academy Sci.*, 5(1):21-23.
- Joshi, N., A. Chauhan and P.C. Joshi. 2009. Impacts of industrial air pollutants on some biochemical parameters and yield in wheat and mustard plants. *The Environmentalist*. 29(4):398-404.
- Lopez, J.M., et al. 2005. Levels of selected metals in ambient air PM₁₀ in an urban site of Zaragoza (Spain). *Env. Res.*, 99(1):58-67.
- Madhavi, L.K. and K.V.S. Badarinath. 2005. Spectral solar attenuation due to aerosol urban area in India. *Atmos. Res.*, 7(4):257-266.
- CPCB. 1995. Implementation status of the pollution control programme in major polluting industries. Programme objectives serives : Probes/62/ 1994-95. Central Pollution Control Board, Delhi.
- Panwar, T.S., D.D.B. Rai and S. Sreekesh. 1997. Ambient air quality status of various cities in India. *UEP*. 17(11):841-845.
- Lakshmi, P.S., K.L. Saravanti and N. Srinivas. 2009. Air pollution tolerance index of various plant species growing in industrial areas. *The Eco. Scan.*, 2(2):203-206.
- Escobedo, F.J., et al. 2008. Analyzing the cost-effectiveness of Santiago Chile's policy of using the urban forest to air quality. *J. Env. Manage.*, 86(1):148-157.
- Lin, Y. J. and H. Ding. 2008. Variation in air pollution tolerance index of plants near a steel factory : Implication for landscape-plant species selection for industrial areas. *WSEAS Transactions Env. Develop.*, 4(1):24-32.
- Ghosh, G.K. 1992. Plants as bio-indicators of automobile exhaust pollution—A case study of Sangi city. *J. Env. Poll.*, 1:26-28.
- Trivedy, M.L. and R.S. Singh. 1995. Reduction in protein contents in few plants as indicators of air pollution. *Poll. Res.*, 14:269-273.
- Pandey, M., et al. 2015. Air pollution tolerance index and anticipated performance index of some plant species for development of urban forest. *Urban Forestry Urban Greening*. 14(4):866-871.

13. Noor, M.J., *et al.* 2015. Estimation of anticipated performance index and air pollution tolerance index and of vegetation around the marble industrial areas of Potwar region : Bio-indicators of plant pollution response. *Env. Geochem. Health.* 37(3):441-455.
14. Nugrahani, P., *et al.* 2012. Ornamental shrubs as plant paletts elements and bio-indicators based on air pollution tolerance index, Surabaya city, Indonesia. *Asian J. Biol. Sci.*, 3(2):295-302.
15. Munshi, A., *et al.* 2001. TRAIL (APO2-L) induces apoptosis in human prostate cancer cells, that is inhabitable by Bcl-2. *Oncogene.*, 20:3757-3765.
16. Sumita, M.N., *et al.* 2003. Tradescantia pallida cv. *Purpurea* boom in the characterization of air pollution by the accumulation of trace elements. *J. Air Waste Manage. Assoc.*, 53(5):574-579.
17. Esfahani, A.A., *et al.* 2013. Assessment of air pollution tolerance index of higher plants suitable for greenbelt development in east of Esfahan city, Iran. *J. Ornamental Horticultural Plants.* 3(2):87-94.
18. Kaur, M. and A. Nagpal. 2017. Evaluation of air pollution tolerance index and anticipated performance index of plants and their application in the development of green space alongwith the urban areas. *Env. Sci. Poll. Res.*, 24(23):18881-18895.
19. Pandey, A. K., *et al.* 2015. Air pollution tolerance index and anticipated performance index of some plant species for the development of the urban forest. *Urban Forestry Urban Greening.* 14(4):866-871.
20. Klumpp, G., *et al.* 2000. Response of stress indicators and growth parameters of *Tibouchina pulchra* Cogn. exposed to air and soil pollution near the industrial complex of Cubatao, Brazil. *Sci. Total Env.*, 246(1):79-91.
21. Flowers, M.D., E.L. Fiscus and K.O. Burkay. 2007. Photosynthesis, chlorophyll fluorescence and yield of snap bean (*Phaseolus vulgaris* L.) genotype differing in sensitivity to ozone. *Env. Exp. Botany.* 61:190-198.
22. Rao, D.N. 1979. Plants as a pollution monitoring device. *Fert. News.* 24:25-28.
23. Hoque, M.A., *et al.* Exogenous proline and glycine betaine increase NaCl-induced ascorbate-glutathione cycle enzyme activities and proline improve salt tolerance more than glycine betaine in tobacco bright yellow-2 suspension-cultured cells. *J. Plant Physiol.*, 164(11):1457-1468.
24. Singh, S.K. and D.N. Rao. 1983. Evaluation of plants for their tolerance to air pollution. Proceedings of Symposium on air pollution control. Indian Association for Air Pollution Control, New Delhi. pp 218-224.
25. Agrawal, S. and V. Tiwari. 1997. Susceptibility level of few plants on the basis of air pollution tolerance index. *Indian Forester.* 123(4):319-322.
26. Dwivedi, A.K. and B.D. Tripathi. 2007. Pollution tolerance and distribution pattern of plants in surrounding area of coal-fired industries. *J. Env. Biol.*, 28(2):257-263.
27. Prajapati, S.K. and B. Tripathi. 2008. Seasonal variation of leaf dust accumulation and pigment content in plant species exposed to urban particulates pollution. *J. Env. Quality.* 37(3):865-870.
28. Govindaraju, M., *et al.* 2011. Identification and evaluation of air pollution tolerant plants around lignite-based thermal power station for greenbelt development. *Env. Sci. Poll. Res. Int.*, 19(4):1210-1223.
29. Shannigrahi, S., T. Fukusima and R.C. Sharma. 2014. Anticipated air pollution tolerance of some plant species considered for green belt development in and around industrial/urban area in India : An overview. *Int. J. Env. Studies.* 61(2): 125-137.
30. Rao, V. L. 2014a. An estimation of air quality indexes of a coastal station-A case study. *Int. J. Current Microbiol. Appl. Sci.*, 3(6):759-763.
31. Rao, V. L. 2014b. Air quality studies over Visakhapatnam-A case study of a coastal station. *Int. J. Innovative Res. Sci. Eng. Tech.*, 1(3): 8215-8220.
32. Kumar, K. S., N. Srinivas and K.A. Sunil. 2014. Monitoring and assessment of air quality with reference to dust particles (PM₁₀ and PM_{2.5}) in urban environment. *Int. J. Res. Tech.*, 3 (16): 42-44.
33. Gajghate, D. G., A.D. Bhanarkar and P. P. Pradeep. 2013. Atmospheric concentration and fluxes of trace elements in Visakhapatnam city of India. *Int. J. Env. Eng.*, 5 (2): 111-128.
34. Gupta, S., *et al.* 2009. Effects of vehicular emissions on biochemical constituents of leaves. *Poll. Res.*, 28:157-160.
35. Agrawal, S.K. 1986. A new distributional function of foliar phenol concentration in the evaluation of plants for their air pollution tolerance index. *Acta Ecol. Sinica.* 8 (2):29-36.
36. Singh, R.P., *et al.* 2007. Biotechnological approaches to improve phytoremediation efficiency for environment contaminants. In *Environmental bioremediation technologies.* Ed S.N. Singh and R.D. Tripathi. Springer, Verlag Berlin Heidelberg. pp 223-258.

37. Prajapati, S.K. 2012. Ecological effect of airborne particulate matter on plants. *Env. Skeptics Critics*. 1(1):12-22.
38. Sen, A., et al. 2017. Eco-physiological evaluation of tree species for bio-monitoring of air quality and identification of air pollution-tolerant species. *Env. Monit. Assess.*, 189(6):262.
39. Dedio, W. 1979. Water relations in wheat leaves as a screening test for drought resistance. *Canadian J. Plant Sci.*, 55:369-378.
40. Barrs, H.D. and P.E. Weatherley. 1962. A re-examination of the relative turgidity technique for estimating water deficit in leaves. *Australian J. Biol. Sci.*, 15:413-428.
41. Lohe, R.N., et al. 2015. A comparative study for air pollution tolerance index of some terrestrial plant species. *Global J. Env. Sci. Manage.*, 4(1):3154-324.
42. Buchchi, B.G., et al. 2013. Evaluation of air pollution tolerance indicate of plant species growing in the vicinity of cement industry and Yogi Vemana University campus. *Indian J. Adv. Chem. Sci.*, 2(1):16-20.
43. Joshi, O.P., K. Pawar and D.K. Wagela. 1993. Air quality monitoring of Indore city with special references to SO₂ and tree barks pH. *J. Env. Biol.*, 14:157-162.
44. Sirajuddin, M., M. Horginamani and M. Ravichandra. 2010. Ambient air quality in an urban area and its effect on plant and human beings : A case study of Tiruchirappali, India. *J. Sci. Eng. Tech.*, 6(2):13-19.
45. Mondal, D., S. Gupta and J.K. Datta. 2011. Anticipated performance index of some tree species considered of green belt development in an urban area. *Int. Res. J. Plant Sci.*, 2(4):99-106.
46. Thambavani, S. and D.V. Prathipa. 2012. Assessment of air quality through bio-monitors of selected sites of Didigul town by air pollution tolerance index approach. *J. Res. Biol.*, 3:193-199.
47. Enete, C., V. Ifeanyi and Ogonna. 2012. Evaluation of air pollution tolerance index (APTII) of some selected ornamental shrubs in Enugu city, Nigeria. *J. Env. Sci. Tech. Food Tech.*, 1(2):22-25.
48. Rao, D.N. and F. LeBlanc. 1996. Effect of sulphur dioxide on the lichen, algae with special reference to chlorophyll. *The Bryologist*. 69(1):89-95.
49. Giri, S., et al. 2013. Effect of air pollution on chlorophyll content of leaves. *Current Agric. Res. J.*, 1(2):93-98.
50. Lima, J.S., E.B. Fernandes and W.N. Fawcett. 2000. *Mangifera indica* and *Phaseolus vulgaris* in the bio-indication of air pollution in Bahia, Brazil. *Ecotoxicol. Env. Safety*. 46:275-278.
51. Chaudhary, C.S. and D.N. Rao. 1997. Study of some factors in plants controlling their susceptibility to sulphur dioxide pollution. *Proceeding Indian National Sci. Academy Part B*. 46:236-24.
52. Garg, A., P. Saxena and C. Ghosh. 2015. Evaluation of tolerance and sensitivity of selected plant species with special references to gasoline exhaust pollution. *Int. J. Sci. Tech. Res.*, 4(2):199-207.
53. Nagendra, H. and D. Gopal. 2010. Street trees in Bangalore: Density, diversity, composition and distribution. *Urban Urban Green*. DOI: 10.1016/j.ufug. 2009.1012.1005.
54. CPWD. 2013. A handbook of landscape. A guide published by Director General. Central Public Works Department, New Delhi.

Scope Of Recycling Of Municipal Wastewater

Debyasmita Sarkar, Sushovan Sarkar* and Debabrata Mazumder

Dr Sudhir Chandra Sur Degree Engineering College, Department of Civil Engineering, Kolkata - 700 074, West Bengal, India

*Corresponding author, Email : hod_ce@dsec.ac.in; dmazumdar@hotmail.com; sdebyasmita@gmail.com

The present research work is done with a goal of achieving a successful way to recycle the municipal wastewater for non-potable daily uses, like toilet flushing, washing, watering in gardens, etc., with a minimum technical knowledge, cost and labour. Grab samples were collected for characterization and performance study on the municipal wastewater. In order to fulfill this objective a suitable channel is selected carrying municipal sewage water from nearby locality. The disposal of such wastewater without any treatment process is going on day by day and thereby creating the alarming situation in surface water quality. The purpose of the present study was to provide a viable treatment option for the sake of recycling. The research was mainly focused on minimum treatment option for municipal wastewater so that the treated effluent could be recycled for non-potable daily uses.

KEYWORDS

Recycling, Municipal wastewater, Non-potable uses, Characterization, Performance study

REFERENCES

1. Devi, M. G., *et al.* 2008. Wastewater reuse and recycling systems : A perspectives into India and Australia. International Water Management Institute. pp 1-44.
2. Jefterson, B., *et al.* 1999. Technologies for domestic wastewater recycling. In Urban water 1. Elsevier Science. pp 285-292.
3. Kamizoulis, G., *et al.* Wastewater recycling and reuse practices in Mediterranean region. Recommended guidelines. pp 1-22.
4. Azarpanah, A. and A. Hajgozar. 2012. Recycle and reuse of wastewater practices in Middle East agriculture. The 1st International and the 4th National Congress on Recycling of organic waste in agriculture. Proceedings, pp 1-6.
5. Asano, T. 2002. Water from wastewater–The dependable water resource. *Water Sci. Tech.*, 48(8): 23-33.
6. Jhansi, S.C. and S.K. Mishra. 2013. Wastewater treatment and reuse : Sustainability options. *J. Sustainable Develop.*, 10(1):1-15.
7. USEPA. Primer for municipal wastewater treatment systems. EPA 832-R-04-001. U.S. Environmental Protection Agency, Office of Water Office of Wastewater Management, Washington, D.C.
8. Morud, J. Reclamation and reuse of wastewater. pp 1-10.
9. Mayer, P.W., *et al.* 1999. Residential end uses of water. American WaterWorks Association Research Foundation, AWWA, USA.
10. Vigneswaran, S. and M. Sundravadivel. Recycle and reuse of domestic wastewater. Encyclopedia of Life Support Systems (EOLSS). pp 1-29.
11. NRC. 1998. Issues in potable reuse–The variability of augmenting drinking water supplies with reclaimed water. National Research Council. National Academy Press, Washington, D.C.
12. IS 10500. 1991. Drinking water specification (1st revision). Bureau of Indian Standards, New Delhi.

Statistical Study Of The Quality Of Water Intended For Irrigation, Case Study Of Region Of Fez, Morocco

M. El Ouchy^{1,2}, K. Arouya^{3,4*}, M. Machkor⁵ and F. Zerrouq²

1. Hospital El Ghassani, Regional Laboratory of Epidemiologic Diagnosis and Hygiene of the Medium, Regional Management of Health, Fez, Morocco

2. High School of Technology, Laboratory of Catalysis, Materials and Environment, Fez, Morocco

3. Ibn Tofail University, Laboratory of Materials, Electrochemistry and Environment, Faculty of Sciences, Kenitra, Morocco

4. Sidi Mohamed Ben Abdellah University, Laboratory of Natural Resources and Environment, Faculty of Polydisciplinary of Taza, Route of Oujda, Taza, Morocco

5. Moulay Ismail University, Faculty of Sciences, Department of Chemistry, Materials, Membranes and Processes of Separation Team, Zitoune, Meknes, Morocco

*Corresponding author, Email : aroyakhalid@gmail.com

This work is part of the characterization of the chemistry of water for irrigation in the region of Fez in Morocco. To carry out this study, 22 samples were taken during two different periods of the year (high water period and rainy return period). Principal component analysis (PCA) applied to the data sets revealed that among the eighteen variables studied, fourteen participate in the constitution of the factor axes. The four remaining variables, namely pH, Ni, Co and As, therefore, do not play an important role in determining the typology of the biotope studied. Moreover, the results obtained indicate that the physico-chemical and toxic (heavy metals) quality of the water used for crop irrigation still does not meet the criteria for use in agriculture. The irrigation water used in the stations located downstream of the industrial discharges of the city of Fez is characterized by high percentages of heavy metals, such as chromium. The protection of water resources and the treatment of polluted water are necessary to ensure that this water can be used in agriculture.

KEYWORDS

Principal component analysis, Heavy metals, Physico-chemical, Water quality, Fez, Morocco

REFERENCES

1. O.M.S. 1989. Use of wastewater in agriculture: Recommendations for sanitary purposes. WHO report 778, Geneva.
2. Bouaicha, R. and A. Benabdelfadel. 2010. Variability and management of surface water in Morocco. *Drought*. 21(1): 1-5.
3. Glaizal, M., et al. 2015. Post-urine assays of heavy metal chelation and pseudoscience: Action must be taken. *Analy. Clinical*. 27(2).
4. Alibou, J. 2002. Impacts of climate change on water resources and wetlands in Morocco. Regional Round Table in the Mediterranean Athens, Greece. pp 1-39.
5. Chahlaoui, A., et al. 1997. Quantitative analysis of the water quality of the Boufekrane wadi, Morocco. *Bulletin Natural History Society*. 133: 71-76.
6. Derwich, E., et al. 2008. Assessment of the quality of surface water from Oueds Fes and Sebou used in market gardening in Morocco. *Larhyss J.*, 7(6): 59-77.
7. A.B.H.S. 2003. Water resources in the Sebou basin, Bilan 2000-2001. Sebou Hydraulic Basin Agency, Fez, Morocco.
8. Benaabidate, L. 2000. Characterization of the Sebou watershed, hydrology, water quality and geochemistry of thermal springs. Thesis in Hydrology at FST- Sais.
9. Azzaoui, S. 2002. Copper, lead and manganese in the Sebou catchment area. *Sources Contribution Impact Quality Surface Water*. 37(4): 733-784.
10. El Ouchy, M., et al. 2016. Analytical validation of method for control of pH and electrical conductivity. *J. Chem. Pharm. Res.*, 8(6):477-484.

11. El Ouchy, M., *et al.* 2014. Confirmation of validation of the method for the determination of nitrite, nitrate and orthophosphate in the water by colorimeter. *J. Mater. Env. Sci.*, 5(S1): 2293-2302.
12. Rodier, J. 1996. The analysis of water (7th edn). **Ed Dunod**. Paris.
13. Rodier, J., L. Bernard and M. Nicole. 2009. Water analysis (9th edn). **Ed Dunod**. Paris.
14. Sampling guide. 2006. Sampling techniques for physico-chemical analyzes. The river quality group led by the Loire-Bretagne Water Agency.
15. Official Bulletin, N ° 5062-30 ramadan 1423.2002. Kingdom of Morocco.
16. Normes and Standards Committee. 1994. Ministry of the Environment of Morocco.
17. Lalami, A. El Ouali. 2011. Surface water pollution in the city of Fez in Morocco: Typology, origin and consequences. *Larhyss J.*, 9(12): 55-72.
18. Derwich, E., *et al.* 2011. Physico-chemical characterization of the waters of the alluvial aquifers of the upper Sebou downstream of its confluence with the Fez wadi. *Larhyss J.*, 8(6): 101-112.
19. Makhoukh, M., *et al.* 2011. Contribution to the physico-chemical study of surface waters of Oued Moulouya. *Larhyss J.*, 9:149-169.
20. Neal, C., *et al.* 2000a. The water quality of the river Kennet: Initial observations on a lowland chalk stream impacted by sewage inputs and phosphorus remediation. *Sci. Total Env.*, 251/252: 477- 495.
21. Moussa, A. B., A. Chahlaoui and R. El Habib. 2013. Change in the physico-chemical quality of Oued Khoumane water in the vicinity of the town of Moulay Idriss Zerhoun, Morocco. *Sci. Lib. J.*, 5.
22. Gouaidia, L. 2008. Influence of lithology and climatic conditions on the variation of the physico-chemical parameters of the water of a water table in semi-arid zone, case of the North-East Algerian Meskiana water table. Doctoral Thesis. Faculty of Earth Sciences, Badji Mokhtar-Annaba University, Algeria.
23. Khamar, M. 2002. Study of the load of metals in water, sediments, soils and agricultural productions irrigated by wastewater of the city of Fez. Doctoral Thesis. Faculty of Science Dhar El Mahraz, Fez.
24. Foudeil, S., *et al.* 2013. Study of heavy metal pollution in Oued Sebou (Morocco). *Sci. Lib. J.*, 5.
25. Chen, W., *et al.* 2013. Accumulation of Cd in agricultural soil under long-term reclaimed water irrigation. *Env. Poll.*, 178: 294-299.
26. Amri, N., *et al.*.2007. Evaluation of the heavy metals accumulate in samples of the sediments, soils and plant by lcpoes with the average Sebou. *J. Condensed Matter*. 8(1).
27. Bouthir, F. Z. 2006. Impact of industrial discharges on the environment. *Mar. Life*. 16:37-47.
28. Koumolou, L., *et al.* 2013. Health-risk market garden production linked to heavy metals in irrigation water in Benin. *Comptes Rendus Biologies*. 336:278–283.

E-Waste Recycling Activities In Global Production And Removal Of Precious And Toxic Metals From Contaminated Water And Soil For Environmental Safety: A Review

Renu Nayar*

D.P. Vipra College, Department of Chemistry, Bilaspur, India

*Corresponding author, Email : nayar.renu@yahoo.co.in

Electronic waste components contain acids, toxic metals and hazardous compounds that may cause health problems and pollute our atmosphere. It is created from anything which is electronic: computers, TV's, monitors, cell phones, vcrs, cd players, fax machines, printers, household equipment's, like vacuum cleaner, microwave/ovens, washing machines, air conditioners, etc., and some smart devices, such as iPhone, iPad, watches, kindles, chromium tablet, etc. Most electronics that are improperly thrown away contain some form of harmful materials, such as lead, copper, cadmium, mercury, beryllium, lithium, barium, nickel, chromium and carcinogenic chemicals which affect the environment and human life. These heavy metals can leach through the soil to reach groundwater which eventually runs to the surface water causing harmful effects to the entire biota either directly or indirectly. Many of these e-waste products can be reused, refurbished, or recycled. Therefore, in the present work, a critical review on environmental effects and removal of toxic metals derived from e-waste recycling activities have been planned. In the present work, the release of these toxic chemicals have been studied and possible remediation processes are discussed.

KEYWORDS

Electrical waste, Toxic metals and compounds, Environment, Recycling activities

REFERENCES

1. Tansel, B. 2017. From electronic consumer products to e-waste : Global outlook, waste quantities, recycling challenges. *Env.Int.*, 98:35-45.
2. Chen, C., *et al.* 2017. Understanding compulsive smartphone use : An empirical test of a flow-based model. *Int. J. Information Manage.*, 37(5):438-454.
3. Dharini, K., *et al.* 2017. IOP Conf. Series. *Earth Env. Sci.*, 80012057.
4. Schmidit, C.W. 2002. E-junk explosion. *Env. Health Perspect.*, 110:A188-A194.
5. Schmidt, C.W. 2006. Unfair trade : E-waste in Africa. *Env. Health Perspect.*, 114: A 232-A 235.
6. Ha, Nguyen Ngoc, *et al.* 2009. Contamination by trace elements at e-waste recycling sites in Bangalore, India. *Chemosphere.* 76:9-15.
7. Pradhan and Kumar. 2014. Informal e-waste recycling : Environmental risk assessment of heavy metal contamination in Mandoli industrial area, Delhi, India. *Env. Sci. Poll. Res. Int.*, 21(13):7913-7928. DOI: 10.1007/s11356-014-2713-2.Epub 2014 Mar 22.
8. Akortia, *et al.* 2017. Soil concentration of polybrominated diphenyl ethers and trace metals from an electronic waste dump site in the Greater Accra region, Ghana. Implications for human exposure. *Ecotoxicol. Env. Safety.* 137:247-255.
9. Fogarasi, S. 2012. Eco-friendly leaching of base metals from waste printed circuit boards : Experimental study and mathematical modeling. In Stud. Univ. Babes-Bolyai Chemia. pp 91-100.
10. Meskers, *et al.* 2009. Impact of pre-processing routes on precious metal recovery from PCs. European metallurgical Conference (EMC). Innsbruck, Austria.
11. Dhawan, N., *et al.* 2008. Recovery of metals from electronic scrap by hydrometallurgical route. Global symposium on Recycling, waste treatment and clean technology (REWAS). Cancun, Mexico. Proceeding, pp 693-698.
12. Wu, K., *et al.* 2015. Heavy metal contamination of soil and water in the vicinity of an abandoned e-waste recycling site : Implications for dissemination of heavy metals.
13. Wong, *et al.* 2007a. Trace metal contamination of sediments in a e-waste processing village in China. *Env. Poll.*, 145:434-442.

14. Wong, *et al.* 2007b. Evidence of excessive release of metals from primitive e-waste processing in Guiyu, China. *Env. Poll.*, 148:62-72.
15. Wong, *et al.* 2007c. Export of toxic chemicals-A review of the case of uncontrolled electronic-waste recycling. *Env. Poll.*, 149:131-140.
16. Leung, *et al.* 2008. Heavy metals concentrations of surface dust from e-waste recycling and its human health implications in southeast China. *Env. Sci. Tech.*, 42:2674-2680. Doi:10.1021/es07187zx.
17. Ha, *et al.* 2009. Contamination by trace elements at e-waste recycling sites in Bangalore, India. *Chemosphere*. 76:9-15.
18. Atiemo, M. National Nuclear Research Institute, Ghana Atomic Energy Commission.
19. Brigden, K., *et al.* 2008. Chemical contamination at e-waste recycling and disposal sites in Acera and Koforidua, Ghana. Greenpeace laboratories technical note. pp 1-23.

River Water Quality Assessment By Bio-Monitoring Working Party Score Of Macro-invertebrate

Janmoni Moran*

Dibrugarh University, Dibrugarh, Assam, India

*Corresponding author, Email : janmonimoran87@gmail.com

An assessment of river water affected by oil refinery effluent was carried out by biological water quality criteria (BWQC) of macroinvertebrate (organisms retained by mesh sizes of ~200-500 mm) community of 2 rivers Dhansiri and Kaliani of Golaghat district of Assam. A total of 35 families of macroinvertebrate were recorded during the study period. The point of effluent discharge showed heavy pollution with the water quality class D and all the control stations of upstream showed water quality class B indicating slight pollution, whereas the downstream of contaminated area of the 2 rivers showed water quality class C indicating moderate pollution.

KEYWORDS

Assam, Community, Effluent, Macroinvertebrate, Pollution, Refinery

REFERENCES

- Allan, J.D. and A.S. Flecker. 1993. Biodiversity conservation in running water. *J. Biosci.*, 43:32-43.
- Schleiger, S.L. 2000. Use of an index of biotic integrity to detect effects of landuses on stream fish communities in west-central Georgia. *Transactions American Fisheries Soc.*, 129:1118-1133.
- Stoddard, J.L., et al. 2006. A process for creating multimetric indices for large-scale aquatic surveys. *J. North American Benthological Soc.*, 27:878-891.
- Beg, M.U., et al. 2003. Distribution of petroleum hydrocarbon in sediment from coastal area receiving industrial effluent in Kuwait. *Ecotoxicol. Env. Safety*. 54:47-55.
- Aghalino, S.O. and B. Eyinla. 2009. Oil exploration and the marine pollution : Evidence from the Niger Delta, Nigeria. *J. Human Ecol.*, 28(3):177-182.
- CPCB Newsletter. 1999. Bio-mapping of rivers. Central Pollution Control Board, New Delhi. 5(4).
- Clements, W.H., et al. 2000. Heavy metals structure benthic communities in Colorado mountain streams. *Ecol. Applications*. 10(2):626-638.
- Fore, L.S. 2002. Biological assessment of mining disturbance on stream invertebrates in mineralized areas of Colorado biological response signature : Indicator patterns using aquatic communities. T.P. Simon. CRC Press LLC, Boca Raton, Florida.
- Miltner, R.J. and E.T. Rankin. 1998. Primary nutrients and the biotic integrity of rivers and streams. *Freshwater Biol.*, 40:145-158.
- Hynes, H.B.N. 1960. The biology of polluted waters. Liverpool University Press, Liverpool.
- Losos, B. 1984. The influence of pollution on the density and production of Chironomidae (Diptera) in running waters. *Limnological (Berlin)*. 15:7-19.
- Armitage, P.D. and J.H. Blackburn. 1985. Chironomidae in a pennine stream system receiving mine drainage and organic enrichment. *Hydrobiologia*. 121:165-172.
- Rao, S.V.R., V.P. Singh and L.P. Mall. 1978. Pollutational studies of river Khan (Indore). I : Biological assessment of pollution. *Water Res.*, 12:555-559.
- Verma, S.R., A.K. Tyagi and R.C. Dalela. 1978. Pollution studies of a few rivers of western U.P. with reference to biological studies. *Proceedings of the Indian Academy Sci.*, 878:123-131.
- Pennak, R.W. 1989. Freshwater invertebrates of the United States : Protozoa to Mollusca (3rd edn). John Willey and Sons, New York, U.S.A.
- Peckarsky, B.L., et al. 1990. Freshwater macro-invertebrates of northeastern North America. Cornell University Press, Ithaca, New York.

17. CPCB Newsletter. 2005. Bio-mapping of rivers-Case study Assam state. Central Pollution Control Board, New Delhi.